

Sociodemographic correlates of older adult acceptance of the COVID-19 vaccine

Arlene Supremo^{1,*} , Sillmark Bacason , Alexander Rex Sañosa 

¹ Biliran Province State University, Philippines

*Correspondence: Arlene Supremo. Address: Biliran Province State University, Philippines. Email: arlene.supremo1986@gmail.com

Responsible Editor: Praba Diyan Rachmawati

Received: 2 September 2021 ◦ Revised: 3 January 2022 ◦ Accepted: 3 February 2022

ABSTRACT

Introduction: An increase of the COVID-19 global statistics in late 2019 prompted the swift manufacturing of vaccines to protect people, especially older adults, from the debilitating effects of the disease. This study aimed to determine the socio-demographic correlates of older adults and their relationship with COVID-19 vaccine acceptance.

Methods: This correlational study invited the entire population of older adults to answer the survey. In total, 89 elderly individuals participated in face-to-face interviews because of the old age limitations. The tool used consisted of two parts. The first part determined the socio-demographic correlates. The second part assessed the level of acceptance of the COVID-19 vaccine. Rank Biserial and Spearman Rho were used to measure the correlations between the socio-demographic variables and COVID-19 vaccine acceptance.

Results: The respondents were predominantly female (n = 45; 50.56%), with an elementary level of education (n = 48; 53.93%), below PHP 10,000.00 for monthly family income (n = 77; 86.52%), unemployed (n = 79; 88.76%), and Roman Catholic (n = 87; 97.75%). Generally, older adults will not accept the COVID-19 vaccine. Specifically, older adults will not accept the vaccine if it has less than a 50% effectiveness (n = 51; 57.30%) and if they have existing medical conditions (n = 51; 57.30%). Finally, the findings suggest that employment status has a significant negative relationship with the acceptance of the COVID-19 vaccine (r = -0.357, p = 0.0001).

Conclusions: 47.19% to 57.30% of older adults will not accept the COVID-19 vaccine. National and local government units need to intensify their vaccination campaigns to persuade the public to engage with the free COVID-19 vaccines.

Keywords: acceptance; COVID-19; hesitancy; pandemic; vaccine; vulnerable

Introduction

The novel coronavirus (COVID-19) pandemic, which started in Wuhan, China, is a global threat, with more than 3 million new reported cases in March 2021 (World Health Organization, 2021). The high statistics in terms of the number of COVID-19 cases prompted nations worldwide to develop mitigating policies to curb the surge of new infections. The health policies include the mandatory wearing of face masks and other personal protective equipment (PPE), as well as isolation and quarantine protocols, granular and community lockdowns, physical distancing, limiting through to prohibiting super spreading interactions like prayer meetings, social gatherings, and personal meetings, and hand sanitization. These non-pharmacological interventions have slowed the rate of coronavirus transmission (El-Elimat et al., 2020). However, pharmacological interventions such as vaccination

remain the best solution to prevent COVID-19 infection (Chakraborty & Parvez, 2020).

Scientists from around the world raced against time to develop potential vaccines against coronavirus. In 2020, the WHO (2021) reported that at least seven vaccines were available for administration. Vaccines are considered helpful in protecting an individual from coronavirus infection through immunity development (CDC, 2021). In the Philippines, the Department of Health (DOH, 2021) conducted an initial vaccine rollout of the Chinese-made SinoVac and UK-made Astrazeneca vaccines to healthcare frontline workers.

Among those who most need the COVID-19 vaccine is the older population group, specifically individuals 65 years old and above, aside from frontline healthcare workers. This group was identified by the WHO (2021) as among the high-risk individuals clustered together with healthcare providers and persons with co-morbidities

because they comprise the majority of the morbidity and mortality cases of the disease. The Centers for Disease and Control Prevention (CDC, [2020](#)) recommended that the older population should be included in the first layer of recipients of the COVID-19 vaccine since the risk of severe illness from the infection increases with age.

With the availability of the COVID-19 vaccine, the next concern that must be overcome by the health authorities and government is its acceptability by older adults as vaccine recipients. Understanding the perspectives of the vaccine recipients is essential when determining the best strategy to maximize vaccine uptake. This milestone in the vaccine roll-out for the older adult population sought to prevent an increase in the disease's morbidity and mortality cases.

Locally, the older adults' acceptance and hesitancy regarding the COVID-19 vaccines have remained unexplored. According to the report by the Biliran Provincial Health Office (BPHO, [2020](#)), there was a low level of compliance with routine immunizations such as the pneumococcal and flu vaccines by older adults. Because of this, a considerable number of older adults were vulnerable to acquiring the infection. This study, therefore, sought to investigate the level of acceptance of COVID-19 vaccination among older adults.

Specifically, this study has been undertaken to determine the sociodemographic profile of older adults in terms of age, sex, education level, monthly income, employment status, and religion. Furthermore, this study wanted to determine the level of acceptance of COVID-19 vaccination among the older adults residing in one of the regions in the Philippines. After determining the sociodemographic profile and level of acceptance of older adults, the causal relationship between the variables was established.

Materials and Methods

Study Design

This study was a correlational research study in terms of its design. Correlational research is concerned with establishing relationships between two or more variables in the same population or between the same variables in two populations. Curtis, Comiskey, and Dempsey ([2016](#)) emphasized that exploring the relationships among the variables is a significant part of research in nursing. Understanding the associations and relationships among human phenomena is a consistent impetus for scientific inquiry in all social science disciplines. This impetus transcends even the most polarized paradigmatic distinctions between various research methods (Fitzgerald et al., [2004](#)). Since the study's goal was to measure and explore the degree of relationship between the demographic characteristics of the older adults and their acceptability of the COVID-19 vaccine, it was appropriate to utilize a correlational research design.

Respondent

The COVID-19 coronavirus pandemic is a global health disaster of the 21st century. This pandemic has caused untold fear and suffering for older adults across the world. According to the World Health Organization (WHO) as cited in the paper by Chee ([2020](#)), older adults in aged care homes are at a higher risk of infection due to living in an enclosed environment with others. Older people are more susceptible to COVID-19 infection, caused by a defective immune response to infectious challenges (Benksim, Addi, & Cherkaoui, [2020](#)). In addition, comorbid illnesses, malnutrition, drugs, and stress predispose the elderly to an increased risk of coronavirus by declining their immune function (Benksim, Addi, & Cherkaoui, [2020](#)). Hence, this study sought the participation of the older adults in the community of Barangay Imelda.

In the Municipality of Naval, Barangay Imelda is held to be one of those barangays with the greatest number of older adults. The barangay is also the barangay furthest away from the urban center of Naval. This confirms their information inaccessibility regarding the benefits of the COVID-19 vaccine. For these reasons, the elderly in this barangay were relevant when it came to providing the data needed for this study.

The researchers used the list of the elderly provided by the barangay health unit as its sampling frame. The elderly listed reached 126 in total. Due to the small population, the researchers used complete enumeration as its sampling technique. However, only 89 were surveyed because only those who were still able to process information were included in the study.

Table 1 Socio-demographic profile of the older adults (n=89)

Demographic Variable	n	%
Sex		
Male	44	49.44%
Female	45	50.56%
Educational Attainment		
Did Not Attend School	2	2.25%
Pre-School Level	3	3.37%
Pre-School Graduate	2	2.25%
Elementary Level	48	53.93%
Elementary Graduate	31	34.83%
Junior High School Level	2	2.25%
College Level	1	1.12%
Family Income		
Below PHP 10,000.00	77	86.52%
PHP 10,000.00 - Php 19,000.00	12	13.48%
Employment Status		
Employed	10	11.24%
Unemployed	79	88.76%
Religion		
Roman Catholic	87	97.75%
Iglesia Filipina Independiente	1	1.12%
Others	1	1.12%

Excluding the few who were not capable of answering the survey increased the reliability of the results.

Instrument

The study developed a questionnaire that would capture its objectives. This self-made questionnaire consisted of 13 items and was pilot tested in Barangay P.S. Eamiguel. The internal consistency of the items was measured using Cronbach's Alpha. The alpha coefficient for the 13 items was 0.992, suggesting that the items had a relatively high internal consistency. To note, a reliability coefficient of 0.70 or higher is considered "acceptable" in most social science research situations. This includes research situations in the field of health and nursing.

The research instrument was divided into two. The first section contained the questions on the demographic profile: (1) Sex, (2) Educational Attainment, (3) Family Income, (4) Employment Status, and (5) Religion of the respondents. The second section contained the 13 items capturing the acceptance of the older adults of the COVID-19 vaccines. The details of the items are presented in [Table 2](#). Furthermore, the respondents rated the variable "acceptance to COVID-19 vaccines" using a five-point Likert scale (Definitely not = 1, Probably not = 2, Probably = 3, Very probably = 4, Definitely = 5).

Data Collection

Initially, the researchers sought ethical clearance from the Research and Innovation Office of the Biliran

Province State University, and permission from the Local Government Unit before the conducting of the study. Informed consent was secured with an emphasis on the study's intention and coverage. Subsequently, the researchers sought assistance from the Barangay Health Office to be provided with a list of older adults in the community. The list was then used as a reference to identify and locate the residences of the older adults in the barangay. Before letting the respondents answer the survey, transmittal letters were given to each of the respondents. After receiving approval through the signing the consent form to survey them, the researcher gave them the survey questionnaire. All of this happened in a face-to-face set-up.

The survey lasted last for a week. A repetition of the barangay visits was done until the data was completed. Since the number of respondents was manageable, the researchers were also those who conducted the survey. They did not seek the help of research assistants or enumerators from the research office of the university.

To emphasize, the survey followed the Inter-Agency Task Force (IATF) guidelines for COVID-19 when visiting the residences of the respondents.

Data Analysis

After gathering the data, the researchers encoded the responses in Microsoft Excel to summarize the data. The responses to some of the variables, particularly those used for profiling, were coded accordingly. Consequently, the researchers produced a descriptive summary with the

Table 2 COVID-19 vaccine acceptance

Statement	Definitely	Very Probably	Probably	Probably Not	Definitely Not
If a vaccine against COVID-19 that has less than 50% effectiveness is available in the market, I will accept it.	9	8	9	12	51
If a vaccine against COVID-19 that has more than 50% effectiveness in the market, I will accept it.	19	5	9	11	45
If a vaccine against COVID-19 has known minor side effect (such as dizziness, lightheadedness, fever, flu-like symptoms (cold, sneezy, body malaise), body pain) and is available on the market, I will accept it.	13	13	7	8	48
I will accept a COVID-19 vaccine if my other family members will take the vaccine shot too.	24	5	9	6	45
I will accept a COVID-19 vaccine if my friends and significant others will take the vaccine shot too.	24	9	3	5	48
I will accept a COVID-19 vaccine if it was recommended by the local health authorities.	26	7	2	8	46
I will accept a COVID-19 vaccine if it was recommended by the national health authorities.	25	9	5	7	43
I will accept a COVID-19 vaccine if it was proven safe and effective by the government.	27	6	6	6	44
I will accept a COVID-19 vaccine if information about the vaccine is provided by trusted health professionals before vaccination starts.	23	11	7	6	42
I will accept a COVID-19 vaccine if information about the vaccine is seen on online platforms such as Facebook, Instagram, etc.	8	12	10	9	50
I will accept a COVID-19 vaccine even if I have existing medical health problems (diabetes, hypertension, rheumatoid arthritis, etc).	13	7	8	10	51
I will accept a COVID-19 vaccine despite my old age.	26	5	2	9	47
I will accept a COVID-19 vaccine even if I am fearful of the vaccination outcome.	21	6	4	8	50

encoded data to determine the completeness of all observations.

The study used SPSS version 25 to analyze and determine the correlation of the demographic variables in relation to the acceptability of the older adults in terms of the COVID-19 vaccine variable. The statistical tools used to measure the correlations of the variables were the Rank Biserial and Spearman Rho. Rank Biserial was used to determine the degree of relationship between the Sex and Acceptance of the older adults of the COVID 19 vaccine and between Employment Status and the Acceptance of the older adults of the COVID 19 vaccine. Meanwhile, Spearman Rho was used to measure the other three demographic variables (Age, Educational Attainment, and Religion) in relation to the Acceptance of the older adults of the COVID 19 vaccine. Sex and Employment Status were run using a different statistical tool of correlation because their data was discrete and nominal in nature. The authors used the p-values of the correlation coefficients to reject or accept the null hypotheses at a significance level of 0.05 and below.

Data Collection

Initially, the researchers sought ethical clearance from the Research and Innovation Office of the Biliran Province State University, and permission from the Local Government Unit before the conducting of the study. Informed consent was secured with an emphasis on the study's intention and coverage. Subsequently, the researchers sought assistance from the Barangay Health Office to be provided with a list of older adults in the community. The list was then used as a reference to identify and locate the residences of the older adults in the barangay. Before letting the respondents answer the survey, transmittal letters were given to each of the respondents. After receiving approval through the signing the consent form to survey them, the researcher gave them the survey questionnaire. All of this happened in a face-to-face set-up.

The survey lasted last for a week. A repetition of the barangay visits was done until the data was completed. Since the number of respondents was manageable, the researchers were also those who conducted the survey. They did not seek the help of research assistants or enumerators from the research office of the university.

To emphasize, the survey followed the Inter-Agency Task Force (IATF) guidelines for COVID-19 when visiting the residences of the respondents.

Ethical Consideration

This study strictly followed the principles of ethical standards when conducting the research. The researchers obtained ethical clearance from the University Internal Review Panel before the actual conducting of the study. Before the data collection process, the researchers provided transmittal letters to the Mayor, Barangay Captain, and Municipal Health Officer.

Three basic principles were observed in the conducting of this study: the principles of respect for persons, beneficence, and justice. Privacy and confidentiality were strictly observed throughout. The respondents in this study were informed of the purpose of the research and provided a clear description of the extent of their involvement and the risks, benefits, and rights of being a study respondent. They were also given the study's strengths related to the actual determination of the acceptance level of older adults for COVID-19 vaccination. However, the actual implementation of the study was constrained due to strict adherence to the COVID-19 protocols, such as the wearing of a face mask, physical distancing, and limited hand hygiene.

A standard form of informed consent was read, elaborated on, and explained in the language that the respondents understood to secure the elements of adequate information and comprehension. The respondents were not coerced, forced, or bribed to participate in this study to secure the element of voluntary participation. The respondents were of legal age and not of unsound mind as declared by the court to secure the element of competency and they qualified based on the inclusion criteria of the study. It was emphasized that they could withdraw from the study at any time during the conducting of the research. The respondents were reminded that they could stop participating in the study at any time without any danger of repercussions.

All data gathered in this study is considered private and confidential. All materials gathered were handled with utmost care. The anonymity of the respondents was preserved. No identifying data was exposed in the report of the study including any names, addresses, etc. All data was destroyed upon the completion and publication of this study.

Compensation or the giving of tokens to respondents poses an ethical dilemma. Nonetheless, this study recognizes the ethical principles of justice and respect and the respondent's time and effort will be given due credit and recognition. Before starting the interview, a token was offered to the respondents. The respondents

Table 3 Employment status and COVID-19 vaccine acceptance

Statement	Correlation Coefficient	p-value	Decision
Q1	-0.414**	0.000	Reject H ₀
Q2	-0.383**	0.000	Reject H ₀
Q3	-0.286**	0.007	Reject H ₀
Q4	-0.315**	0.003	Reject H ₀
Q5	-0.305**	0.004	Reject H ₀
Q6	-0.317**	0.003	Reject H ₀
Q7	-0.311**	0.003	Reject H ₀
Q8	-0.288**	0.006	Reject H ₀
Q9	-0.317**	0.002	Reject H ₀
Q10	-0.342**	0.001	Reject H ₀
Q11	-0.394**	0.000	Reject H ₀
Q12	-0.344**	0.001	Reject H ₀
Q13	-0.357**	0.001	Reject H ₀

*<0.05 **<0.01

were informed that the interview tokens were given based on their assessed needs.

Lastly, there was no known potential conflict of interest for this study.

Results

Out of the 89 respondents who participated in the study, 45 (50.56%) were female and 44 (49.44%) were male. Most of the respondents had reached the elementary level of education, totaling 48 (53.93%). Concerning family income, many older adults belong to a family that receives a monthly revenue of less than PHP 10,000.00 per month as most of them are unemployed (88.76%). Roman Catholicism (97.75%) is the predominant religion of the group.

The older adults will definitely not accept the vaccine in the circumstances presented in [Table 2](#). However, more older adults will definitely not accept the vaccine if it has less than 50% effectivity and if they have existing medical conditions such as diabetes, hypertension, and rheumatoid arthritis at 51% and 57%, respectively.

Regarding understanding the correlation between the socio-demographic correlates and the acceptance of the COVID-19 vaccine, only the employment status of older adults, particularly those who were unemployed, showed a significant correlation. The majority of the socio-demographic correlates (sex, educational attainment, family income, and religion) had a negative correlation with COVID-19 vaccine acceptance.

Discussions

The COVID-19 pandemic has wreaked havoc on the global economy since its first occurrence in Wuhan, China. Because of the negative pandemic effects, the WHO and its partners, especially drug manufacturers, have raced against time to develop a vaccine that will help reduce the statistics of COVID-19 infection (WHO, [2021](#)). The available COVID-19 vaccines ready for administration include SinoVac Astrazeneca, Moderna, Sputnik V, and Pfizer. However, successful vaccines will still rely on the uptake of the population (Zigron et al., [2021](#)).

As of July 7, 2021, the WHO ([2021](#)) reported that 3,032,056,355 people have received the COVID-19 vaccine out of the 7,753,000,000 total population around the world (World Bank Group, [2021](#)). In the Philippines, the COVID-19 vaccination rollout began in March 2021 to eligible priority groups which included older adults (Department of Health, [2021](#)). The acceptance of these older adults of the available COVID-19 vaccines has been studied here in relation to their sociodemographic profile.

The COVID-19 vaccine acceptance of the older adults showed that most people belonging to this population group will definitely not accept the vaccine. Among the notifiable response where most older adults did not show

acceptance of the COVID-19 vaccine was if the vaccines had less than 50% effectiveness. Harapan et al. ([2020](#)) and Alqudeimat et al. ([2021](#)) described in their studies that acceptance is relative to the effectiveness of a certain COVID-19 vaccine. The higher the effectiveness, the more people will definitely accept the vaccine. Therefore, the government must put out effective strategies that will persuade the population to avail the free COVID-19 vaccines if the one purchased has been known to have low effectiveness.

Furthermore, older adults who have existing comorbidities such as diabetes and hypertension expressed their unwillingness to accept the COVID-19 vaccine. The previous literature showed a similar finding that people with underlying medical conditions are hesitant and resistant to accepting the COVID-19 vaccines (Murphy, [2021](#)). Unacceptance was linked to the medical contraindication of vaccines and the fear of vaccine side effects. This result is counterintuitive to the somatic benefit-risk analysis where people with health problems should take extra protective measures from infectious diseases. Older adults with existing health problems are considered vulnerable to COVID-19 infection (CDC, [2021](#)), therefore they need the vaccine more. The unacceptance of the COVID-19 vaccine from this group might hamper the achievement of herd immunity where more people are expected to be fully vaccinated.

Upon studying the relationship between the sociodemographic correlates of the older adults in relation to their willingness to accept the COVID-19 vaccine, only unemployment showed a significant correlation. This information revealed a contrasting result to those of the available studies in the literature where unemployed individuals are less likely to accept the COVID-19 vaccines (Malik et al., [2020](#); Echoru et al., [2021](#)). Malik et al. ([2020](#)) reported that vaccine acceptance is directly proportional to literacy level. The higher the education level, the more accepting an individual is of the COVID-19 vaccine.

Strikingly, the study shared the same result as the study by El-Elimat et al. ([2020](#)). Their study reported that unemployed participants were more accepting of the COVID-19 vaccines than the employed participants. This is not a perplexing finding as most of the respondents in the study were unemployed due to old age. Most of them were retired individuals receiving pensions. Others were supported by their family.

Limitations in relation to the conducting of this study were identified. One limitation was the research locale, wherein all study participants lived in a mountainous barangay. A different finding may have been gathered if the study encompassed a wider coverage of older adults – those living in rural-upland, rural-coastal, and urban areas, respectively. Also, the study was conducted when COVID-19 vaccines were still undergoing clinical trial. As more studies proving the safety and effectiveness of the

COVID-19 vaccines emerge, the respondents might change their perception regarding acceptance.

Conclusions

Generally, the results of this study highlight the need to improve the level of knowledge and increase the trust in the COVID-19 vaccines to expedite their uptake in the older adult population. Educating them on the benefits of the COVID-19 vaccine is an essential milestone in preventing further morbidity and mortality cases of the COVID-19 disease, considering their high-risk category. Moreover, the local government should intensify its public awareness campaigns to reach everyone in the community. Providing rewards or incentives can also be a strategy to improve the level of vaccine uptake.

References

- Al-Qerem, W. et al. (2021). 'COVID-19 vaccination acceptance and its associated factors among a Middle Eastern population'. *Frontiers in Public Health*, 9, 34. Retrieved from <https://doi.org/10.3389/fpubh.2021.632914>.
- Alqudeimat, Y. et al. (2021). 'Acceptance of a COVID-19 vaccine and its related determinants among the general adult population in Kuwait'. *Medical Principles and Practice*, 30(3), 262-271.
- Baldacchino, G. (2006). 'Islands, island studies, Island Studies Journal; *Island Studies Journal*, 1(1), 3-18. Retrieved March 29, 2021, from <https://islandstudies.ca/sites/vre2.upei.ca/islandstudies.ca/files/u2/SJ-1-1-2006-Baldacchino-pp3-18.pdf>
- Benksim, A., et al. (2020). 'Vulnerability and Fragility Expose Older Adults to the Potential Dangers of COVID-19 Pandemic'. *Iranian Journal of Public Health*, 49(1), 122-124. Retrieved March 29, 2021, from https://www.researchgate.net/publication/340984493_Vulnerability_and_Fragility_Expose_Older_Adults_to_the_Potential_Dangers_of_COVID-19_Pandemic
- Biliran Provincial Health Office. (2020). *Needs assessment survey of Barangay Imelda Naval*. BPHO: Naval. Biliran.
- Burns, A. (1999). 'Collaborative Action Research for English Language Teachers'. Cambridge University Press.
- Centers for Disease Control and Prevention. (2021). *What older adults need to know about COVID-19 vaccines*. Retrieved from <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/recommendations/older-adults.html>
- Centers for Disease Control and Prevention. (2021). *Benefits of getting a COVID-19 vaccine*. Retrieved from <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/vaccine-benefits.html>.
- Centers for Disease Control and Prevention. (2021). *COVID-19: people with certain medical conditions*. Retrieved from <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>.
- Chakraborty, R. et al. (2020). 'COVID-19: an overview of the current pharmacological interventions, vaccines, and clinical trials'. *Biochemical pharmacology*, 114184. <https://doi.org/10.1016/j.bcp.2020.114184>
- Chee, S. Y. (2020). 'COVID-19 Pandemic: The Lived Experiences of Older Adults in Aged Care Homes'. *Millennial Asia*, 11(3), 299-317. DOI: 10.1177/0976399620958326
- Curtis, E. et al. (2016). 'Importance and use of correlational research'. *Nurse Researcher*, 23(6), 20-25. doi:10.7748/nr.2016.e1382
- Department of Health. (2021). *When will the COVID-19 vaccines be available to me?* Retrieved from <https://doh.gov.ph/Vaccines/when-will-the-COVID-19-available-to-me>
- Echoru, I. et al. (2021). 'Sociodemographic factors associated with acceptance of COVID-19 vaccine and clinical trials in Uganda: a cross-sectional study in western Uganda'. *BMC public health*, 21(1), 1-8. <https://doi.org/10.1186/s12889-021-11197-7>
- El-Elimat, T. et al. (2020). 'Acceptance and Attitudes Toward COVID-19 Vaccines: A Cross-Sectional Study from Jordan'. *medRxiv*. <https://doi.org/10.1101/2020.12.22.20248676>
- Fitzgerald, S. M. et al. (2004). 'Correlational designs in rehabilitation research'. *Journal of Vocational Rehabilitation*, 20(2), 143-150.
- Hair, J. F. et al. (2010). 'Multivariate Data Analysis (7 ed.)'. Pearson Prentice Hall.
- Harapan, H. et al. (2020). 'Acceptance of a COVID-19 vaccine in southeast Asia: A cross-sectional study in Indonesia'. *Frontiers in public health*, 8. doi: 10.3389/fpubh.2020.00381
- Malik, A. A. et al. (2020). 'Determinants of COVID-19 vaccine acceptance in the US'. *EClinicalMedicine*, 26, 100495. DOI: [10.1016/j.eclinm.2020.100495](https://doi.org/10.1016/j.eclinm.2020.100495)
- Murphy, J. et al. (2021). 'Psychological characteristics associated with COVID-19 vaccine hesitancy and resistance in Ireland and the United Kingdom'. *Nature communications*, 12(1), 1-15. <https://doi.org/10.1038/s41467-020-20226-9>
- World Bank Group. (2021). *Population, total*. Retrieved from <https://data.worldbank.org/indicator/SP.POP.TOTL>.
- World Health Organization. (2021). *COVID-19 vaccines*. Retrieved from <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/covid-19-vaccines>
- World Health Organization. (2021). *Weekly epidemiological update on COVID-19-16 March 2021*. Retrieved from <https://www.who.int/publications/m/item/weekly-epidemiological-update---16-march-2021>
- World Health Organization. (2020). *Coronavirus disease (COVID-19): vaccines*. Retrieved from [https://www.who.int/news-room/q-a-detail/coronavirus-disease-\(covid-19\)-vaccines?adgroupsurvey={adgroupsurvey}&gclid=CjwKCAjwOGCBhAlEiwA7FUXkugtjuY2FBIHXby_DO9t0RSrwTDZDc008BtToDzINnh0_4Umn-ytBoCkIQOAvD_BwE](https://www.who.int/news-room/q-a-detail/coronavirus-disease-(covid-19)-vaccines?adgroupsurvey={adgroupsurvey}&gclid=CjwKCAjwOGCBhAlEiwA7FUXkugtjuY2FBIHXby_DO9t0RSrwTDZDc008BtToDzINnh0_4Umn-ytBoCkIQOAvD_BwE)
- Zigron, A. et al. (2021). 'COVID-19 vaccine acceptance among dental professionals based on employment status during the pandemic'. *Frontiers in Medicine*, 8, 13. DOI: [10.3389/fmed.2021.618403](https://doi.org/10.3389/fmed.2021.618403).

How to cite this article: Supremo, A., Bacason, S., and Sañosa, A. A. (2022) 'Sociodemographic correlates of older adult acceptance of the covid-19 vaccine', *Jurnal Ners*, 17(1), pp. 2-7. doi: <http://dx.doi.org/10.20473/jn.v16i1.29655>

The effect of basic trauma and cardiac life support training in increasing the competence of emergency room nurses

Anisha Calista Prakoeswa^{1,*} , Fitri Arofiati¹, Nur Hidayah¹ 

¹ Muhammadiyah Yogyakarta University, Indonesia

*Correspondence: Anisha Calista Prakoeswa. Address: Brawijaya Street, Kasihan District, Bantul Regency, Yogyakarta 55183, Indonesia. Email: anisha.calista.psc20@mail.umy.ac.id

Responsible Editor: Ferry Efendi

Received: 17 February 2022 ◦ Revised: 3 March 2022 ◦ Accepted: 23 March 2022

ABSTRACT

Introduction: Nurses on duty in the emergency room (ER) play an important role in the early identification of a life-threatening patient condition, immediate action, and the subsequent care of critically ill patients. Continuous training and improvement need to be done to improve the ER nurses' competency. This research aimed to analyze the differences in nurses' competencies.

Methods: This quantitative study employed a quasi-experiment design. The sample size comprised 50 nurses of ER in Surabaya with a total sampling technique. Variables used on this study were demographical data as the independent variables and dependent variables including knowledge, attitude and skills. Intervention was given by two methods, online training for knowledge and attitude, while offline training for skill improvement. After ensuring the distribution and homogeneity of the data, a paired t-test was employed for parametric statistical data analysis.

Results: There were differences in pre-test and post-test scores in the domain of knowledge, attitude and skill ($p = 0.000$), respectively. The mean of knowledge, attitude, and skills had increased significantly. Hence, the training is proven to improve those competencies as well as the sub-topic items.

Conclusions: Basic Trauma and Cardiac Life Support (BTCLS) training has improved nurse competency; it is advised to hold this training, especially in the ER. Indeed, it can improve healthcare services quality and improve patient safety at health facility. In addition, future research can modify the training time, training delivery methods, and content of BTCLS training materials to adjust with the current situation of the COVID-19 pandemic.

Keywords: attitude; competency; emergency nurse; knowledge; skill; training

Introduction

The nurse plays a crucial role in determining health services in hospitals, not only because nurses work on the frontline, but also being a majority compared to other health workers (WHO, 2020). By implementing professional nursing care, good quality health services can be accomplished. Nurses who work in the emergency room (ER) must adopt critical situations and provide nursing care with speed and accuracy that is different with other nursing situations (Laoh and Rako, 2014). Nurses on duty in the ER play a vital role in the early identification of a life-threatening patient condition, immediate action, and the subsequent care of critically ill patients (Cunningham *et al.*, 2017). Compared to other patient care units, the ER and ICU are places where the

number of critically injured patients is the most, so it is essential to ensure that nurses working in these units have adequate competence (Sankar *et al.*, 2013).

Nurses must perform continuous self-development by increasing knowledge, attitudes, and skills to provide proper emergency services. Training is the effort to ensure that nurses have competency in emergency cases. This method is an option that does not require much cost but can provide the expected effect even if it is carried out by utilizing limited resources (Cunningham *et al.*, 2017). The competence of nurses in handling emergency cases is essential to increase the success rate, accuracy, and quality of emergency service in treating patients (Mason *et al.*, 2005). Therefore, competency in handling emergency cases is one of the most important factors determining healthcare service quality (Li *et al.*, 2016).

Corona Virus Disease 2019 (COVID-19) is a global crisis that happened in late 2019 and rapidly spread all over the world (Yie *et al.*, 2021). In this constraint, nurses are included in the first line in hospitals that provide healthcare services (Buchan *et al.*, 2019). Therefore, a nurse must have sufficient understanding in recognizing COVID-19 disease, including identifying the pathogen, signs and symptoms, how it spreads, how to break the chain of infection, and the treatment that must be given appropriately (Burnett, 2018; Corless *et al.*, 2018). Nurses' competency regarding COVID-19 disease will contribute to their environment because their competence will provide health protection for themselves and the surrounding community. The efforts to improve the nurses' competency are an investment during the COVID-19 pandemic (Ridley, Sanderson and Haines, 2021). This effort not only provides positive outcomes in dealing with the COVID-19 pandemic but also prepares for another outbreak that may occur in the future (Purba, 2020). To maximize healthcare personnel's capabilities and increase protection, vigilance, and control against COVID-19 infection, hospitals should provide training for nurses, either on-site or online (Wu *et al.*, 2020). The online training was an effective method for improving the competency of nursing workforce either in academic or clinical setting (Wong and Greenhalgh, 2013; Tobase *et al.*, 2017; Vaona *et al.*, 2018). Dr. Soetomo Hospital Surabaya routinely conducted emergency nursing training once a year and was designed for nurses to handle emergency problems. This training used an appropriate approach by referring to scientific foundations and good nursing processes. Hence, the training participant is expected to be able to demonstrate life saving skills, minimize organ damage, and reduce patient mortality and disability. This training was provided to new nurses who will be transferred or assigned to the ER. Competence is a set of abilities for individuals to carry out a job correctly and consists of knowledge, attitudes, and skills. However, research regarding the effect of the training using online and on-site approach toward nurse's competency has not been done.

This study aimed to analyze the differences in nurse competencies, in terms of knowledge, attitude, skills, in ER after attending BTCLS training.

Materials and Methods

Study Design

This quantitative research employed a quasi-experimental study, using a one-group pre-test-post-test design, with BTCLS training as an intervention. BTCLS training was held on 12-16 September 2021 and combining both offline and online sessions to reduce physical interaction during the COVID-19 pandemic. Before the training, the first observations (pre-test)

measured the participants' competencies (knowledge, skills, and attitudes). Then the second observation (post-test) was made again after the training by re-measuring the participants' competencies (knowledge, skills, and attitudes). The first three days of BTCLS training were conducted online. The first day of BTCLS training includes participant registration, technical meeting, health protocol presentation as well as knowledge competency pre-test, while the second and third day of BTCLS training consisted of online lecture materials that includes the role of emergency nurses in improving quality profession during the pandemic, Basic Life Support, trauma of thorax and abdomen, triage, electrocardiogram (ECG) and acute coronary syndromes, burns, and fluid therapy. All of the online sessions of BTCLS training were held by Zoom Meetings. The fourth and fifth day of BTCLS training were conducted offline while maintaining health protocols. All of the participants and instructors took antigen swab test before the offline sessions of BTCLS training. Participants were divided into small groups following the six stations namely basic life support skills, initial assessment, transport skills, airway and breathing skills, ECG skills, and advanced life support skills. On the fourth day, the instructors held the pre-test of both the attitudes and skills of the participants during each station before giving out the demonstrations of the right methods for attitudes and skills. And lastly, the participants went through post-test for knowledge, attitudes, and skills on the fifth day.

Respondent

The sample size comprised 50 nurses with a total sampling technique. The participants worked in ER Dr. Soetomo Hospital Surabaya. The independent variable in this study was BTCLS training. The dependent variable is nurses' competence, which consists of knowledge, skills, and attitudes.

Instrument

Instruments for assessing knowledge, skills, and attitudes were obtained from the official assessment, Education and Training Board of the Regional Supervisory of the Indonesian National Nurses Association (PPNI) East Java; hence there was no need to test for validity and reliability (Zúñiga *et al.*, 2016; Stevanin *et al.*, 2017). The knowledge assessment consists of 35 multiple choice questions according to the material contained in the BTCLS training. Meanwhile, attitude and skill measurement are shaped as a checklist for the participant's emergency care completeness, and the BTCLS training assessors assessed it. The assessors were competent assessor whose having field experience on BTCLS and has been recognized as certified assessor in this area. Each assessment for knowledge, skill, and attitude is considered good if the total score is $> 75\%$ and not good if the score is $< 75\%$.

Table 1 BTCLS Training Participants Based on Age, Gender, Education, and Length of Work

Characteristics	n	%
Gender		
Male	26	52
Female	24	48
Age		
25 – 30 years	11	22
31 – 35 years	27	54
36 – 40 years	7	14
> 40 years	5	10
Unit		
1st floor ER	25	50
Contagious ER	7	14
ER-ROI	2	4
Aster HCU	4	8
HCU Pandan II	1	2
Others	11	22
Education Level		
Bachelor	21	42
Associate Degree (D3)	27	54
Associate Degree (D4)	2	4
Period of Employment		
< 1 year	11	22
1-5 Year(s)	8	16
> 5 Years	31	62

BTCLS training is a nursing training to handle emergency problems caused by trauma and cardiovascular disorders. The training aims to enable nurses to provide life support to save lives and minimize patients' disability and organ damage. Nurse competency is the ability of each nurse to handle emergency cases under training standards. The competency consists of (1) knowledge: what nurses know about handling emergency cases; (2) attitude: how nurses handle emergency patients; (3) skill: nurses' ability to handle

emergency patients according to applicable standards and procedures. Knowledge competency data are nominal. Meanwhile, the measurement scale for skill and attitude is ordinal data.

Data Collection

This research was divided into three stages, namely the preparation stage, implementation, and the final stage. The preparation stage is a preliminary study. The researcher conducted interviews with the Director of Medical & Nursing Services, the Director of Professional Education & Research, the Head of ED's Nurse, and the Head of the Nursing Division. Interviews were conducted to overview existing training and evaluation of emergency nursing throughout this period. Then further discussions were held with representatives from the Indonesian Emergency and Disaster Nurses Association (HIPGABI) to determine emergency nursing training tailored to the COVID-19 pandemic situation.

Data Analysis

The researcher identified the participants and conducted preliminary observations (pre-test). Furthermore, recording and documentation were accomplished during BTCLS training. The final observation (post-test) was conducted to see the effect of the training. In addition, the researcher also conducted additional interviews with the BTCLS training organizers. The final stage was data analysis using the SPSS program. The researcher first looked at the distribution and homogeneity of the data. A paired t-test was employed for parametric statistical data analysis.




Figure 1. Boxplot graph of nurses' competency before and after BTCLS training

Ethical Consideration

In addition, at the implementation stage, the researcher also obtained ethical clearance from the health research ethics committee, Dr. Soetomo Surabaya Hospital. This research was conducted with ethical principles: informed consent, anonymity, and confidentiality. This study received ethical clearance from the ethics commission, with letter number 0240/KEPK/VII/2021, valid for one year since first issued on August 23rd, 2021.

Results

The normality test of the three competencies shows that the data obtained are normally distributed. Kolmogorov-Smirnov test results for pre-test knowledge $p = 0.200$, post-test knowledge $p = 0.200$, pre-test attitude $p = 0.116$, post-test attitude $p = 0.200$, pre-test skill $p = 0.107$, and post-test skill $p = 0.094$.

The homogeneity and normality test are assumptions of the hypothesis test formula for the t-test and ANOVA. Before testing the data by t-test and ANOVA, the assumptions of normality and homogeneity must be met. The next test is the homogeneity test, which determines the data variance. Levene's test results are as follows, knowledge $p = 0.199$, attitude $p = 0.127$, and skill $p =$

Table 2 Results of the pre- and post-data analysis training of BTCLS of nurses in Dr. Soetomo Hospital Surabaya

Variable	Δ Mean	Δ Std. Deviation	t	p-value
Knowledge (pre- and post-test data)	12.940	4.181	21.881	0.000
Attitude (pre- and post-test data)	14.300	5.273	19.176	0.000
Skill (pre- and post-test data)	14.380	5.102	19.928	0.000

Table 3 Results of the pre- and post-data analysis competencies in detail BTCLS training of nurses in Dr. Soetomo Hospital Surabaya

Variable	Δ Mean	Δ Std. Deviation	t	p-value
Knowledge				
Trauma	0.127	0.175	5.144	0.000
Life Support	0.154	0.187	5.844	0.000
Cardiac	0.313	0.210	10.540	0.000
Attitude				
Orientation phase	0.458	0.447	7.240	0.000
Show a listening attitude	0.197	0.368	3.775	0.000
Provide an opportunity to ask questions or clarify	0.700	1.015	4.876	0.000
Documents according to standard	0.760	1.333	4.030	0.000
Termination phase	0.907	0.771	8.316	0.000
Wash your hands according to standard	1.060	1.163	6.443	0.000
Aseptic and antiseptic principles for sterile procedures	1.100	2.092	3.718	0.001
Protection with universal precautions	1.100	2.092	3.718	0.001
Handling medical and non-medical waste	0.500	1.515	2.333	0.024
Skills				
Preparation	0.850	0.649	9.264	0.000
Assessing Patients	0.820	0.569	10.185	0.000
Airway	0.825	0.552	10.577	0.000
Breathing	0.845	0.463	12.916	0.000
Circulation	0.830	0.519	11.316	0.000
Disability	0.713	0.522	9.669	0.000
Exposure	0.750	0.672	7.892	0.000
Foley Catheter	0.927	0.592	11.068	0.000
Gastric Tube	0.873	0.466	13.251	0.000
Heart Monitor	0.760	0.797	6.743	0.000
Pulse Oximetry, X-Ray	0.520	0.707	5.202	0.000
Secondary Survey	0.667	0.738	6.390	0.000

0.088. The three competencies have the same variance, and the assumption of homogeneity has been met.

[Table 2](#) show the results of the BTCLS training effect on nurse competencies (knowledge, attitude, and skills). In column 'p,' the effect is marked with a significance value below 0.05. There are differences in the pre-test and post-test scores of the nurse competency test. Each competency has a significance value of less than 0.000. Hence, this quasi-experimental research hypothesis concludes that BTCLS training significantly affects nurses' knowledge, attitude, and skills. The mean of knowledge has increased significantly by 12.94. This pattern also occurs in other nurse competencies: attitude and skills. The results show that the post-test scores of the majority of nurses are improved. The boxplot graph explains an increase in the post-test scores on knowledge, attitude, and skills compared to the pre-test scores ([Figure 1](#)).

[Table 3](#) shows differences in post-test knowledge and pre-test knowledge in trauma, life support, and cardiac. Each of these categories has a p-value below alpha (0.05). Furthermore, the table also shows the difference of post-test attitude and pre-test attitude in the categories of orientation, listening, an opportunity to ask questions or clarify, standardized documents, termination, hand washing, aseptic and antiseptic principles, protection, and handling of medical and non-medical waste. Finally, the table describes the differences in each category of questions on skill competence. All categories of skill competency questions, including airway, breathing, circulation, disability, exposure, foley catheter, gastric tube, heart monitor, pulse oximetry and x-ray, and secondary survey showed significant differences

Discussions

This research showed that there were significant differences between knowledge before and after training. This finding aligns with Ha and Nuntaboot's (2020) research on nursing education and training as predictors of nurse competence. However, the difference is that the research results emphasize on narrative views' of participants on the training (Ha and Nuntaboot, 2020). This finding also parallels with prior research conducted by Fong *et al.* (2021). It is more relevant since the same methodology -quasi-experimental without a control group- was employed to examine the effect of orientation on nurse competence especially focus on increased knowledge (Fong *et al.*, 2021). Item analysis was also accomplished in this study, with good results in knowledge (trauma, life support, and cardiac). There was a significant difference in the percentage of trainees who answered correctly. Structured training is required in order to increase the competency of health workers in health facilities, including nurses.

This research demonstrated that there were significant differences between attitudes before and after

training. This finding supports prior research on emergency, that training affects the knowledge and skills of healthcare personnel (Yildiz, Selimen and Dogan, 2014; Ameh *et al.*, 2016). However, the difference is that those studies only examined partial competencies (Yildiz, Selimen and Dogan, 2014; Ameh *et al.*, 2016). This study also included item analysis, which yielded positive findings in terms of attitude. There was a significant difference in the percentage of trainees who behave correctly in the orientation phase, listening attitude, opportunity to ask questions/clarification for patients, documents according to standards, termination phase, washing hands, aseptic principles and antiseptic, universal precaution protection, and handling medical and non-medical waste. Future training in increasing attitude should consider the general and specific competency of nurses.

This research revealed that there were significant differences between skills before and after training. This finding confirms prior research conducted by Kim and Shin (2016) in the area of maternal and child emergency nursing. Overall, this research on improving the emergency nurses' competency supports research from Ojifinni, Motara and Laher (2019) to include basic life support (BLS) and cardiopulmonary resuscitation (CPR) in the formal training to improve nurse competency. Park and Kim (2017) also put forward the same suggestion to conduct specific education and training regarding an emergency training to improve the nurses' competency. In this study, item analysis was also performed, and the results were positive for nurses' skill competency. There was a highlighted finding for nurses who acted correctly on preparation, assessing patients, airway, breathing, circulation, disability, exposure, foley catheter, gastric tube, heart monitor, pulse oximetry, x-ray, and secondary survey. Skill of nurses should be viewed as critical competency as it is related to patient safety and improving the quality of healthcare.

This study has several limitations, including without using a control group; since this research is the first to be conducted to assess the increase in knowledge, attitude, skills, and a category of questions at the research locus. Hence, it is hoped that further research can be conducted with re-assessments within a certain period (three months, six months, or one year) so that training can be refreshed to adjust with the needs of nurses.

Conclusions

There was an increase in emergency nurses' competency in term of knowledge, attitude, and skill after attending the BTCLS training. Repeated educational programs can improve these three domains as well as the contents of the domain. For future BTCLS training, recommendations include training time, training delivery methods, and content of BTCLS training materials. In addition, future research can modify the

training method as an intervention group and add a control group to the standard training method.

References

- Ameh, C.A. *et al.* (2016) 'Knowledge and skills of healthcare providers in sub-Saharan Africa and Asia before and after competency-based training in emergency obstetric and early Newborn Care', *PLoS ONE*, 11(12), pp. 1–14. doi:10.1371/journal.pone.0167270.
- Buchan, J. *et al.* (2019) *A critical moment: NHS staffing trends, retention and attrition*, The Health Foundation.
- Burnett, E. (2018) 'Effective infection prevention and control: the nurse's role', *Nursing Standard* [Preprint]. doi:10.7748/ns.2018.e11171.
- Corless, I.B. *et al.* (2018) 'Expanding nursing's role in responding to global pandemics 5/14/2018', *Nursing Outlook*, 66, pp. 412–415.
- Cunningham, C. *et al.* (2017) 'Developing an emergency nursing short course in Tanzania', *African Journal of Emergency Medicine*, 7(4), pp. 147–150. doi:10.1016/j.afjem.2017.08.002.
- Fong, H.C. *et al.* (2021) 'A Competency-Based Orientation (CBO) Protocol Enhances Competency Among Newly Hired Perianesthesia Nurses', *Journal of Perianesthesia Nursing*, 36(5), pp. 445–451. doi:10.1016/j.jopan.2020.09.017.
- Ha, D.T. and Nuntaboot, K. (2020) 'Factors influencing competency development of nurses as perceived by stakeholders in vietnam', *Belitung Nursing Journal*, 6(4), pp. 103–110. doi:10.33546/BNJ.1119.
- Laoh, J. and Rako, K. (2014) 'Gambaran pengetahuan perawat pelaksana dalam penanganan pasien gawat darurat di ruangan IGD BLU RSUP. Prof. dr. r. d kandou manado', *Jurnal Ilmiah Perawat Manado*, 3(2), p. 92833.
- Li, S.-M. *et al.* (2016) 'Research progress in disaster nursing competency framework of nurses in China', *Chinese Nursing Research*, 3(4), pp. 154–157. doi:10.1016/j.cnre.2016.11.003.
- Mason, S. *et al.* (2005) 'Developing assessment of Emergency Nurse Practitioner competence - A pilot study', *Journal of Advanced Nursing*, 50(4), pp. 425–432. doi:10.1111/j.1365-2648.2005.03408.x.
- Ojifinni, K., Motara, F. and Laher, A.E. (2019) 'Knowledge, Attitudes and Perceptions Regarding Basic Life Support Among Teachers in Training', *Cureus*, 11(12). doi:10.7759/cureus.6302.
- Park, H. and Kim, J. (2017) 'Factors influencing disaster nursing core competencies of emergency nurses', *Applied Nursing Research*, 37, pp. 1–5.
- Purba, A.K. (2020) 'How should the role of the nurse change in response to Covid-19?', *Nursing times*, 116(6), pp. 25–28.
- Ridley, E.J., Sanderson, A.F. and Haines, K.J. (2021) 'Surge capacity for critical care specialised allied health professionals in Australia during COVID-19', *Australian Critical Care*, 34, pp. 191–193.
- Sankar, J. *et al.* (2013) 'Knowledge and skill retention of in-service versus preservice nursing professionals following an informal training program in pediatric cardiopulmonary resuscitation: A repeated-measures quasiexperimental study', *BioMed Research International*, 2013. doi:10.1155/2013/403415.
- Stevanin, S. *et al.* (2017) 'The Multidimensional Nursing Generations Questionnaire: development, reliability, and validity assessments', *Journal of Nursing Management*, 25(4), pp. 287–296. doi:10.1111/jonm.12465.
- Tobase, L. *et al.* (2017) 'The effects of an online basic life support course on undergraduate nursing students' learning', *International journal of medical education*, 8, pp. 309–313. doi:10.5116/ijme.5985.cbce.
- Vaona, A. *et al.* (2018) 'E-learning for health professionals', *The Cochrane database of systematic reviews*, 1(1), pp. CD011736–CD011736. doi:10.1002/14651858.CD011736.pub2.
- WHO (2020) *State of the world's nursing 2020: investing in education, jobs and leadership*. Geneva: World Health Organization.
- Wong, G. and Greenhalgh, T. (2013) *Internet-based Learning for Training Health Care Professionals In-service*. London: Queen Mary University of London.
- Wu, X. *et al.* (2020) 'Contingency nursing management in designated hospitals during COVID-19 outbreak', *Annals of Global Health*, 86(1), pp. 1–5. doi:10.5334/aogh.2918.
- Yie, K.Y. *et al.* (2021) 'Using social network analysis to identify spatiotemporal spread patterns of covid-19 around the world: Online dashboard development', *International Journal of Environmental Research and Public Health*, 18(5), pp. 1–15. doi:10.3390/ijerph18052461.
- Yildiz, T., Selimen, D. and Dogan, D. (2014) 'A Study Comparing the Pre- and Post-Training Knowledge of Emergency Department Nurses in Turkey for the Diagnosis of Physically Abused Women', *Journal of Family Violence*, 29(5), pp. 519–525. doi:10.1007/s10896-014-9608-3.
- Zúñiga, F. *et al.* (2016) 'Evidence on the validity and reliability of the German, French and Italian nursing home version of the Basel Extent of Rationing of Nursing Care instrument', *Journal of Advanced Nursing*, 72(8), pp. 1948–1963. doi:10.1111/jan.12975.

How to cite this article: Prakoeswa, A. C., Arofiati, F., and Hidayah, N. (2022) 'The effect of basic trauma and cardiac life support training in increasing the competence of emergency room nurses', *Jurnal Ners*, 17(1), pp. 8–13. doi: <http://dx.doi.org/10.20473/jn.v17i1.33754>

Indonesians' readiness in facing long-term COVID-19 pandemic

Yuyun Setyorini ¹, Yopi Harwinanda Ardesa ² , Rendi Editya Darmawan ^{1,*} 

¹ Department of Nursing, Health Polytechnics of Surakarta, Indonesia

² Department of Orthotics-Prosthetics, Health Polytechnics of Surakarta, Indonesia

*Correspondence: Rendi Editya Darmawan. Address: Letjen Sutoyo Street, Mojosongo, Surakarta, Central of Java, Indonesia,

Email: rendiedityad@gmail.com

Responsible Editor: Yulis Setiya Dewi

Received: 29 July 2021 ◦ Revised: 22 February 2022 ◦ Accepted: 1 March 2022

ABSTRACT

Introduction: Long-term COVID-19 pandemic which has existed in Indonesia since 2020 not only poses a risk to public health but also to their social conditions. Indonesia, as a country with strong social fabric within its people, has a high sense of social responsibilities in facing the pandemic. The aim of this study is to describe Indonesians' readiness in facing long-term COVID-19 pandemic.

Methods: This descriptive study used a cross-sectional design with a sample of 305 residents of five cities in Indonesia. They were chosen not randomized by distributing a survey invitation via WhatsApp group with the background of the general public, and health cadres. We waited until the response received reached the targeted number of respondents. The data were collected using Google Forms, and were analyzed by using a descriptive method (percentage, P-value, and odds ratio) by SPSS statistical software version 25.

Results: The readiness in facing the pandemic fell into two categories: the high readiness level (77.1%) and the low readiness level (22.9%). Gender and age do not affect the level of readiness, while the level of education and employment status influenced the level of readiness.

Conclusions: The majority of Indonesians have high readiness level. Public participation can decrease COVID-19 transmission rate.

Keywords: COVID-19; Indonesian; pandemic; readiness

Introduction

Coronavirus Disease 2019 (COVID-19) is a new strain that has never been identified before in humans (Tarantola, 2020). Based on the scientific facts available during the formulation of the research, COVID-19 can be transmitted human-to-human through droplets of cough or sneeze and is not airborne. Persons with the most risk of contracting this disease are those coming into close contact with COVID-19 patients, including those treating them (Heymann, 2020; Song and Lu, 2020). Standard recommendations in preventing the transmission of the virus include washing hands regularly with soap and running clear water, applying coughing and sneezing etiquette, avoiding direct contact with livestock and wild animals, and avoiding close contact with anyone exhibiting respiratory disease symptoms such as coughing and sneezing (WHO, 2020a; 2020c).

The public needs to be ready with the new habits like maintaining distance, washing hands, wearing masks, and not gathering. On the other hand, Indonesians have a habit of gathering and socializing with family, and friends, and this has the potential for transmission of COVID-19. Another problem is not meeting with family, and friends (Pasteruk, 2020). Not socializing in a short time doesn't cause problems, but not being able to meet for a long time will cause boredom (Hwang et al., 2020). A long-lasting pandemic can make people tired of wearing masks because it's uncomfortable and costs money (Claresta, Christian and Sa'id, 2021). We need to see Indonesian readiness to solve the problem, and carry out health protocols for a long time. Readiness is defined as a point of maturity to accept and practice the health protocol behavior that has been set in life.

The case of COVID-19 has spread throughout the world, numbering 237 countries based on WHO data in

March 2022. The incidence of COVID-19 based on WHO data on March 17, 2022 showed 462,758,117 confirmed cases with a mortality rate of 6,056,725 deaths. The WHO also reported data on confirmed cases of COVID-19 in South-East Asia of 56,626,996 (WHO, 2022). Data from the Indonesian Ministry of Health reported that, as of March 17, 2022, there were 5,939,082 Indonesians confirmed COVID-19 with 153,212 deaths. Based on WHO data in February 2022, Indonesia was placed 27 of 237 countries with the most COVID-19 cases (Kementerian Kesehatan Republik Indonesia, 2022). Jansen et al. (2021) stated that there is a faster transmission in the Omicron variant, making the potential for COVID transmission increase when public awareness is low.

Indonesia's position was included in countries with a high number of COVID-19 cases, in line with data monitoring health protocol compliance in July 2021. Task force COVID-19 reported that 27.03% people did not obey wearing masks, and disobedience to keep a distance was 28.38%. It is not only the responsibility of the government, but also community contribution to increase public participation of the importance for obeying health protocols (Satgas, 2021).

Indonesians have a successful history in solving problems by involving the community, such as a successful program to prevent dengue fever by establishing larva monitoring (Widyawati, 2019). The country also succeeded in controlling drug dropout in tuberculosis patients using drug monitoring, where support from the supervisor significantly influences the patients' obedience of taking TB drugs (Widani and Sianturi, 2020). These problems were solved by the community. History shows that the role of the community is very important in controlling COVID-19 in Indonesia.

Based on community roles win line with public health condition, community readiness in carrying out, inviting, and monitoring health protocols becomes important. Community readiness depends on individual commitment to change, facilitating change, and implementing change (Edwards et al., 2000; Castañeda et al., 2012). Indonesians are accustomed to the same behavior as their values. For instance, the residents of Sukoharjo, Province Central Java decided to isolate the village as a form of war against the coronavirus, and no one is allowed in or out until the situation return to normal, and each individual follows the rules (Pasteruk, 2020).

Seeing the long-term likelihood of COVID-19 pandemic, as well as compliance with fluctuating health protocols, a study was needed to determine the readiness of Indonesians in dealing with a lengthy COVID-19 pandemic. There was no research that discusses Indonesians' readiness in deal with a long-term COVID-19 pandemic, hence the objective of this research was to investigate Indonesians' readiness in facing a long-term COVID-19 pandemic.

Materials and Methods

Study Design

This research used a descriptive study with a cross-sectional design and was carried out in Indonesia between March and June 2021.

Respondent

All participants (n=305) were Indonesians living in five cities: Surakarta, Surabaya, Jayapura, Denpasar, and Sukoharjo. The data of the research were collected through questionnaire via Google Forms including demographic information (age, sex, level of education, and employment status), and measures of readiness in coping with COVID-19 undertaken.

Instrument

The measures of readiness were assessed by using the 5-point Likert scale to measure responses for 14 questions. The instrument made by the research team in consultation with linguists, and public health expert was then tested for validity and reliability. Validity and reliability tests were carried out on 20 respondents outside the research sample. Validity test was using Pearson product moment and reliability test using Cronbach's alpha. The results of the r table with 20 respondents and the level of significance is 5%, namely 0.444. A statement is declared valid if r count > r table. The questions 1 through 14 have r count more than r table and it can be concluded that all statements are valid. The results of Cronbach's alpha show a value of 0.786 and are more than the r table (0.444), so it can be concluded that all statements are reliable (Appendix 1).

Table 1 Demographic characteristics and readiness level of the respondents (n = 305)

Variables	Frequency	Percentage
Gender		
Female	266	87.2%
Male	39	12.8%
Age (years)		
17-35	69	22.6%
36-50	157	51.5%
>50	79	25.9%
Education level		
Elementary school	20	6.6%
Junior high school	45	14.8
Senior high school	139	45.6
Diploma Degree	35	11.5
Bachelor's degree	42	13.8
Master's degree	24	7.7
Occupation		
Unemployed	175	57.4%
Civil servants	35	11.5
Traders	19	6.2
Laborers	16	5.2
Students	21	6.9
Others	39	12.8
Readiness Level		
High readiness level	235	77.2%
Low readiness level	70	22.8%

Table 2 Relationship between characteristics of the respondents and readiness level

Variable	Readiness level		OR (95% CI)	P value*
	High (%)	Low(%)		
Gender				
Female	208 (68%)	58 (19%)	1.237	0.591
Male	29 (10%)	10 (3%)		
Age				
17-50 years old	171 (56%)	55 (18%)	0.612	0.150
>50 years old	66 (22%)	13 (4%)		
Level of education				
High	78 (26%)	32 (10%)	1.739	0.000
Low	157 (51%)	38 (12%)		
Employment status				
Employed	71 (23%)	26 (9%)	2.164	0.000
Unemployed	166 (54%)	42 (14%)		

Data Collection

We distributed survey invitations via WhatsApp to WhatsApp groups with the background of the general public, and health cadres, and waited until the response received reached the targeted number of respondents.

Data Analysis

The results show that Indonesians' readiness in facing the pandemic fell into two categories, namely the high readiness level and the low readiness level. Readiness level was determined by identifying 14 statements ([Appendix 2](#)). The statement consists of readiness in identifying guests after the trip, quarantine for people with a high risk of COVID-19, checking the health of residents, reducing the frequency of mobility, providing supporting facilities for health protocols, disinfection of public places, health education, participation in surveillance of suspected COVID-19, and sharing food. Each statement will get a value of 1-5. The maximum value for each respondent is 70. The result of the calculation of mean is 53. We measured group readiness levels with a formula, readiness value more than or equal to 53 was referred to as high readiness level, and readiness value less than 53 as low readiness level. The readiness measures collected were then grouped into those that carried the most (high readiness level) and those carried the least (low readiness level). Data analysis was performed by using the SPSS statistical software version 25 by showing percentage, odds ratio (OR), and P-value. The data were presented or summarized as mean, frequency, OR, and percentage.

Ethical Consideration

The study was approved by the Research Ethics Committee of the Indonesian Ministry of Health. The Ethical Approval was obtained from Health Research Ethics Committee of Poltekkes Kemenkes Surakarta, number LB.02.02/1.1/2424.4/2021 dated on January 31st, 2021. Permission to conduct the study was proposed to the Regional Research and Development Design Agency of Central Java Province. The researchers applied research ethics principles of anonymity, beneficence and non-maleficence, autonomy, and justice. An explanation of the background and aim of the study, as well as informed

consent were contained in the Google Form, together with the instrument. Participants were reminded at the beginning of the survey that proceeding and completing the survey indicated voluntary consent to participate in the study. Anonymity and confidentiality were maintained throughout the study.

Results

A total of 305 people was sampled with 95% response rate of 321 respondents who were recruited. The respondents ranged from 17-63 years of age SD of 34±6.3 years. [Table 1](#) shows demographic data of the respondents. The majority of participants were female (87.2%). A large number of respondents were within the age range of 36-50. Most of respondents (45.6%) graduated from Senior Secondary School, and a large number of respondents (57.4%) were unemployed. The majority of respondents had a high readiness level (77.1%).

Based on the research, it is reported that women (68%) had higher level of readiness in facing COVID-19. Respondents falling into productive age of 17–50 were readier in dealing with the impacts of the pandemic. It is also reported that respondents with higher levels of education were 1.739 times readier than those with lower levels of education, and those who were employed were 2.164 times readier than those who were not ([Table 2](#)).

Discussions

The research results show that the majority of Indonesians (77.1%) were of high readiness level in facing COVID-19. The factors influencing readiness in facing the pandemic were education level ($p = 0.000$) and employment status ($p = 0.000$), while gender ($p = 0.591$) and age ($p = 0.150$) were not found to be correlated with public readiness level. This result is influenced by several determinants of readiness.

The respondents with higher levels of education had 1.739 times higher readiness level than those with lower levels of education. Education and skills have a strong influence on individual well-being. Education creates many opportunities for individuals and is essential for high economic growth, and strong social cohesion

(Mojtahedi et al., 2021). Education guarantees income, social benefits, healthy neighbors, and healthy behaviors, so that those with higher levels of education are readier to face new diseases. This is in line with findings from Yanti et al. (2020) that show Indonesians have good knowledge, positive attitudes, and good behavior toward social distancing to prevent the transmission of the COVID-19 pandemic in Indonesia. It can be concluded that level of knowledge influences public readiness in facing COVID-19.

The respondents who were employed were 2.164 times readier than those who were unemployed. This is consistent with the findings of Weil (2017), stating that employment influences one's level of readiness in facing problems. Employed people will have better economic state, which enables them to make ends meet. Mojtahedi et al. (2021) said that people who lost their jobs during the pandemic felt depression, anxiety, and stress, so they had difficulty facing COVID-19.

As a consequence of the COVID-19 pandemic, many companies terminated their staff's employments. The rules of social distancing have forced coffee shops to lay off their employees. The implementation of the lockdown has also prevented the tourism, hotel and transportation sectors from allowing employees to work (Al-Fadly, 2020; Botton, Hoffmann and Vera-Cossio, 2020). If this global health crisis is not resolved immediately, the economy will not do well, thereby causing an increase in vulnerability of transmission and rate of death (Viezzer and Biondi, 2021).

The results of the study show that most of the men and women have high readiness. Level of exposure risk, comorbidity, socioeconomic level, and individual experiences influence readiness in facing the pandemic. The WHO recommends that, in making decisions, governments should also consider gender factors in order to ensure public participation and readiness (WHO, 2020b).

Age correlates with level of experience and vulnerability to disease. In this research it was found that those aged 17-50 years, as well as those more than 50 years old, were in high readiness to face COVID-19, although Indonesians had no experience in dealing with diseases similar to COVID-19. Experience in facing diseases similar to COVID-19 provides more readiness. As an example, Taiwanese, as reported by Chiu (2020) were ready to face COVID-19 since they had experience in facing SARS in 2003. The researchers stated that COVID-19 prevention efforts have been conducted earlier in Taiwan, including countering false news spreading fear (Chiu, 2020). Vulnerability to COVID-19 should cause people to be more vigilant, in which the elderly are more vulnerable to the disease (Niu et al., 2020). However, since Indonesians have no prior experience, age did not significantly influence readiness.

Indonesians voluntarily raised funds to make washbasins, distribute face masks, disinfect places of

worship, and help others economically since they have a culture of mutual assistance (*gotong royong*), which has long taken root in the country (Bowen, 1986). The history of three centuries of colonization enables Indonesians to achieve high levels of solidarity. Cooperation, mutual assistance, collective decision-making, and mutual respect are important parts in the national culture (Dewantara, 2017). The limitation of this research is that the sample coverage is not comprehensive as it does not cover all 34 provinces in Indonesia.

Conclusions

Research findings show that the majority of Indonesians sampled have high readiness level in facing the COVID-19 pandemic. Most males and females in Indonesia, both aged 17-50 years or more than 50 years, who are highly educated and have low education, as well as those who are working or not working have high readiness to face the pandemic. There is, however, a need of improvement in health education in order to improve readiness. We suggest that a wider survey is needed by adding a variable about COVID-19 vaccination. Health protocol compliance must also always be surveyed as a basis for determining government policies.

References

- Al-Fadly, A. (2020) 'Impact of Covid-19 on SMES and Employment', *Entrepreneurship and Sustainability Issues*, 8(2), pp. 629–648. doi: 10.9770/jesi.2020.8.2(38).
- Botton, N., Hoffmann, B. and Vera-Cossio, D. (2020) 'The unequal impact of the coronavirus pandemic: Evidence from seventeen developing countries', *PLoS ONE*, 15(10), pp. 1–10. doi: 10.1371/journal.pone.0239797.
- Bowen, J. R. (1986) 'On the Political Construction of Tradition: Gotong Royong in Indonesia', *The Journal of Asian Studies*. [Cambridge University Press, Association for Asian Studies], 45(3), pp. 545–561. doi: 10.2307/2056530.
- Castañeda, S. F. et al. (2012) 'Dimensions of community and organizational readiness for change', *Progress in Community Health Partnerships: Research, Education, and Action*, 6(2), pp. 219–226. doi: 10.1353/cpr.2012.0016.
- Chiu, W. T. (2020) 'Determinants of Taiwan's Early Containment of COVID-19 Incidence', *AJPH*, 110(7), pp. 943–945. doi: 10.2105/AJPH.2020.305720.
- Claresta, A., Christian, S. and Sa'id, M. (2021) 'Public Obedience to Health Protocols during Covid-19 Pandemic in Indonesia: A Perspective from Health Belief Model Theory', *Journal of Social Sciences and Humanities*, 11(2), pp. 1–11.
- Dewantara, A. (2017) "'Gotong-Royong' (Mutual Assistance of Indonesia) According to Soekarno in Max Scheler's Axiology Perspective', *International Journal of Humanities and Social Science*, 4, pp. 41–50. doi: 10.14445/23942703/IJHSS-V4I5P106.
- Edwards, R. W. et al. (2000) 'The Community Readiness Model: Research to Practice Ruth', *Journal of Community Psychology*, 28(3), pp. 291–307. doi: 10.1002/(SICI)1520-6629(200005)28.
- Heymann, D. L. (2020) 'A novel coronavirus outbreak of global health concern', *thelancet*, 395, pp. 15–18. doi: 10.1016/S0140-6736(20)30185-9.
- Hwang, T. J. et al. (2020) 'Loneliness and social isolation during the COVID-19 pandemic', *International Psychogeriatrics*, 32(10), pp. 1217–1220. doi: 10.1017/S1041610220000988.
- Jansen, L. et al. (2021) 'Investigation of a SARS-CoV-2 B.1.1.529 (Omicron) Variant Cluster — Nebraska, November–December 2021', *MMWR. Morbidity and Mortality Weekly Report*, 70(5152), pp. 1782–1784. doi: 10.15585/mmwr.mm705152e3.
- Kementerian Kesehatan Republik Indonesia (2022) *Situasi Covid-19 Nasional*.
- Mojtahedi, D. et al. (2021) 'The Relationship Between Mental Toughness, Job Loss, and Mental Health Issues During the

COVID-19 Pandemic', *Frontiers in Psychiatry*, 11(February), pp. 1–16. doi:10.3389/fpsy.2020.607246.

Niu, S. *et al.* (2020) 'Clinical characteristics of older patients infected with COVID-19: A descriptive study.', *Archives of Gerontology & Geriatrics*. Beijing Emergency Medical Center, Beijing, China: Elsevier B.V., 89, p. N.PAG-N.PAG. doi: 10.1016/j.archger.2020.104058.

Pasteruk, I. (2020) 'Community Development in Indonesia: Contemporary Aspects of Culture', *Proceedings of the International Conference on Community Development*, 477(Icccd), pp. 338–342. doi: 10.2991/assehr.k.201017.075.

SatgasCovid-19 (2021) *Monitoring Pemantauan Protokol Kesehatan di Wilayah Indonesia, Satuan Tugas Penanganan COVID-19*.

Song, F. and Lu, H. (2020) 'Emerging 2019 Novel Coronavirus (2019-nCoV) Pneumonia', *Radiology*, 295(1), pp. 210–217.

Tarantola, D. (2020) 'A Public Health Perspective in the Times of COVID-19', *AJPH*, 110(7), pp. 925–927. doi: 10.2105/AJPH.2020.305742.

Viezzler, J. and Biondi, D. (2021) 'The influence of urban, socio-economic, and eco-environmental aspects on COVID-19 cases, deaths and mortality: A multi-city case in the Atlantic Forest, Brazil', *Sustainable Cities and Society*, 69(June), pp. 1–15. doi: <https://doi.org/10.1016/j.scs.2021.102859>.

Weil, A. R. (2017) 'The Work/Health Relationship', *Health Affairs*, p. 199. doi: 10.1377/hlthaff.2017.0059.

WHO (2020a) *Advice on the use of masks the community, during home care and in health care settings in the context of the novel coronavirus (2019-nCoV) outbreak Interim guidance*.

WHO (2020b) *Gender and COVID-19*.

WHO (2020c) *Novel Corona Virus, WHO*.

WHO (2022) *WHO Coronavirus (COVID-19) Dashboard, WHO*.

Widani, N. L. and Sianturi, S. R. (2020) 'Relationship Between Drug Consumption, Supervisors' Knowledge and Support, and Patients' Obedience to Take Tuberculosis Drugs', *IJNP (Indonesian Journal of Nursing Practices)*, 4(1), pp. 46–52. doi: 10.18196/ijnp.41107.

Widyawati, W. (2019) *Satu Rumah Satu Jumantik Efektif Cegah DBD, Kementerian Kesehatan*.

Yanti, B. *et al.* (2020) 'Community Knowledge, Attitudes, and Behavior Towards Social Distancing Policy As Prevention Transmission of Covid-19 in Indonesia', *Jurnal Administrasi Kesehatan Indonesia*, 8(2), pp. 4–14. doi: 10.20473/jaki.v8i2.2020.4-14.

How to cite this article: Setyorini, Y., Ardesa, Y. H., and Darmawan, R. E. (2022) 'Indonesians' readiness in facing long-term covid-19 pandemic', *Jurnal Ners*, 17(1), pp. 14–18. doi: <http://dx.doi.org/10.20473/jn.v17i1.28707>

Appendices

Appendix 1 Validity and reliability readiness level assessment

Statements	Pearson r Count	Cronbach's alpha
Statement 1	0.715	0.786
Statement 2	0.617	
Statement 3	0.694	
Statement 4	0.568	
Statement 5	0.782	
Statement 6	0.600	
Statement 7	0.785	
Statement 8	0.671	
Statement 9	0.598	
Statement 10	0.638	
Statement 11	0.641	
Statement 12	0.723	
Statement 13	0.652	
Statement 14	0.568	

Appendix 2 Readiness level assessment

Statements	Totally Agree (5)	Agree (4)	Partially Agree (3)	Disagree (2)	Totally Disagree (1)
1) Do you carry out identity checks and ask the city of origin where the migrants come from?					
2) Do you ask visitors to check with the nearest health facility after the trip?					
3) Do you ask visitors to apply self-quarantine measures for 14 days inside their homes after the trip?					
4) Do you ask the arriving person to self-quarantine at the quarantine facility provided by the city government?					
5) Do you check the body temperature and health status of the people who come?					
6) Do you do road closures to reduce the frequency of mobility?					
7) Do you build a sink and provide soap?					
8) Do you distribute masks to the public?					
9) Do you disinfect places of worship and public places?					
10) Do you make health education posters or banners?					
11) Do you suspend meetings and rallies?					
12) Do you raise funds to ease the economic burden of the population?					
13) Do you participate in surveillance activities against suspected COVID-19?					
14) Do you donate food to members of the community affected by COVID-19 who are in need					

**Scoring: Total score: _____

The association between self-control and satisfaction in life among Filipino older adults

Laarni A. Caorong ^{1,*} 

¹ Mindanao State University, Philippines

*Correspondence: Laarni A. Caorong. Address: Main Campus, Marawi City, 9700 Lanao del Sur, Philippines, Email: laarni.caorong@msumain.edu.ph

Responsible Editor: Rizki Fitriyarsari

Received: 7 March 2022 ◦ Revised: 10 April 2022 ◦ Accepted: 12 April 2022

ABSTRACT

Introduction: Self-control has been associated with a host of positive life outcomes. This correlational study was aimed at determining the link between self-control and life satisfaction among the older population.

Methods: This study used a quantitative correlational design. Ninety-six (96) older adult residents of Iligan City and Lanao del Norte, Mindanao Philippines, who met the inclusion criteria were recruited. This quantitative study made use of two standardized instruments namely the Brief Self-Control Scale (BSCS) and the Satisfaction with Life Scale (SWLS). The association was assessed using Pearson correlation statistics.

Results: There is a significant relationship between the older adults' self-control and their life satisfaction. The computed *r* value is 0.248 ($p < 0.05$). Evidently, there is a positive relationship between self-control and life satisfaction. This current study has provided evidence to support the assumption that exercising self-control leads to life satisfaction.

Conclusions: As a significant positive relationship between the older adults' self-control exercise and their life satisfaction was proven, this means that effective exercise of self-control positively leads to better life outcomes such as satisfaction in life. Linking self-control exercise with life satisfaction offers an innovative way for us to know why some people may feel more satisfied with life than others.

Keywords: elderly; Filipino; life satisfaction; older adults; self-control

Introduction

The concept of attaining life satisfaction has always been an aspiration to most human beings. While self-control is attributed to a host of positive outcomes, it is still interesting to know the link between self-control and the satisfaction with life. The essence of self-control in attaining important life outcomes is widely recognized. High self-control, according to several authors, positively predicts a person's well-being, life satisfaction and positive affect. High self-control here means the ability of a person to regulate his thoughts, feelings and behaviors (De Ridder et al., 2012; Hofmann et al., 2014).

Several authors have pointed out that self-control is a powerful determinant of success across the life span (De Ridder et al., 2012). The practice of self-control is associated with positive outcomes such as better interpersonal relationships, good physical health and better intellectual performance (Finkel and Campbell, 2001; Schmeichel, Vohs and Baumeister, 2018). In this study, self-control is defined as the self's capacity to

override or change one's inner responses, as well as to interrupt undesired behavioral tendencies and to refrain from acting on them (Tangney, Boone and Baumeister, 2018).

Exercising high self-control is a pertinent aspect of a person's behavior for a person to have a successful and healthy life (De Ridder et al., 2012; Tangney, Boone and Baumeister, 2018). It was found that behaviors such as minimized aggression, reduced criminality improved relationships, less abuse of alcohol, smoking and other prohibited substances and high self-esteem and improved interpersonal skills were all related to the practice of high self-control (Finkel and Campbell, 2001; Sayette and Griffin, 2004). Moreover, Hofmann et al. (2014) stated that self-control is a person's ability to override one's inner response and to interrupt undesirable behavioral inclinations or impulses.

Additionally, greater self-control is also positively attributed to psychological adjustments and negatively predicts psychopathology (Tangney, Boone and



Baumeister, 2018). It was also found that the more self-control a person exhibits, the fewer are the experience of symptoms and stress while having a better mental health (Boals, Vandellen and Banks, 2011). This finding was corroborated with the outcome of the study that there is a link between self-control and higher quality and satisfying relationships (Bogg and Roberts, 2004; Jensen-Campbell and Malcolm, 2007).

Moreover, several researchers have perceived self-control as a key aspect in personal conscientiousness, which is subsequently linked to longevity, physical health and other relevant health behaviors (Friedman et al., 1993; Bogg and Roberts, 2004; Goodwin and Friedman, 2006). Succinctly, self-control practice is an all-encompassing important human quality trait essential for attaining a good life, thus its practice results in many positive life outcomes.

The aging population is rapidly growing worldwide, and attention must be given to the older sector of the society. There are certain conditions associated with aging such as decline in economic resources, decrease in cognitive ability, failing health and weakening social support (Deary et al., 2009). The various changes that take place in old age suggest that aging might be linked to a decline of the older adults' well-being. An important measure of subjective well-being is life satisfaction. Life satisfaction is established as an essential indicator of wellness in general. It is also seen as a necessary gauge of positive functioning (Gilman, 2009). Numerous people and several researchers find getting old linked to a decline in happiness in older people. This is attributed to the physical, social and psychological changes that they experience in old age which clearly exhibit that the experiences that older people go through impact their satisfaction with life.

The theory of self-control in old age has an assumption that the exercise of self-control is linked to life satisfaction (Caorong, 2019). The aim of this study is to know and determine if the exercise of self-control is associated with life satisfaction. Determining the association between self-control and life satisfaction will help confirm the assumption of the theory of self-control in old age which states that the exercise of self-control among older adults leads to life satisfaction. The result of this study will give a better perspective on the concept of life satisfaction in late life and will provide and enhance the viewpoint on the link between life satisfaction and self-control.

Materials and Methods

Study Design

This study utilized a quantitative correlational research design to determine the relationship between the self-control of older people and their life satisfaction.

Respondent

The target participants were 60 years old and older adults from Iligan City and Lanao del Norte, the Philippines. Target population (N=96) was calculated based on correlation sample size calculator [$\alpha=0.05$, $\beta=0.50$, $r=0.2$]. Participants were purposively chosen to answer the research instruments. Inclusion criteria included: being a male or female, sixty years old and above, and those older adults demonstrating normal cognitive and mental functioning. Cognitive functioning was assessed utilizing the Short Portable Mini-mental Status Questionnaire or SPMSQ by Pfeiffer (1975). Participants who committed two or less errors were asked to join the study. However, those who committed three or more errors using the (SPMSQ) indicating cognitive impairment were not included.

Instrument

The Brief Self-Control Scale (BSCS) is a standardized instrument which measures self-control. It is comprised of statements which are measured on a scale ranging from 1 to 5 where 1 indicates 'not at all like me' and 5 indicates 'very much like me' (Tangney, Boone and Baumeister, 2018). The scale consists of a total of thirteen items with an internal consistency estimate (alpha) of 0.83 and 0.85 in two study samples. The test-retest reliability estimate was 0.87 (three-week interval). Self-control as defined by Tangney is the self's capacity to override or change one's inner responses, as well as to interrupt undesired behavioral tendencies and to refrain from acting on them (Tangney, Boone and Baumeister, 2018).

The Satisfaction with Life Scale (SWLS) is another standardized instrument used in this study which measures the life satisfaction element of subjective well-being. Life satisfaction judgments represent the cognitive component of subjective well-being, or the experience commonly referred to as happiness (Diener et al., 1985, 1999). The SWLS is Likert in type where a seven-point response category is used for the 5-item statements comprising the scale. The lowest score is five while the highest possible score that can be obtained is 35. Score interpretation included the following: 5-9 (extreme dissatisfaction with life); 10-14 (dissatisfied with life); 15-19 (slightly dissatisfied with life); 20 (neutral); 21-25 (slightly satisfied with life); 26-30 (satisfied with life) and 31-35 (extremely satisfied with life). The scale has a high internal consistency with a coefficient alpha of 0.79 to 0.89. Additionally, in a month interval, the test-retest correlation scores were 0.84 and 0.80.

Data Collection

The researcher collected data by distributing a standardized research instrument for the participants to answer. The instrument comprised the Brief Self-control Scale developed by Tangney, Baumeister and Boone (2018) and the Satisfaction with Life Scale (SWLS) by

Table 1 Profile of the participants (N = 96)

Variable	n	%
Age		
60-69 years old	60	62.50
70-79 years old	31	32.29
80 years old and above	5	5.20
Marital Status		
Married	51	53.12
Widow	37	38.54
Single	3	3.12
Separated	5	5.20
Religion		
Roman Catholic	82	85.41
Islam	9	9.37
Others	5	5.20
Educational Attainment		
No Formal education	18	18.75
Elementary Level	30	31.25
Elementary Graduate	7	7.29
High School Level	12	12.50
High School Graduate	8	8.33
College Level	7	7.29
College Graduate	10	10.41
Postgraduate Level	4	4.16
Employment Status		
Unemployed	53	55.20
Employed	43	44.7
Monthly Income (in PHP)		
5,000 and below	76	79.16
5,000-9,999	7	7.29
10,000-14,999	7	7.29
15,000-19,999	2	2.08
20,000-24,999	1	1.11
25,000-29,999	1	1.11
30,000 and above	2	2.08

Table 2 Descriptive statistics for the older adults' level of self-control (N = 96)

Self-control Categories	n	%
Very low (13-26)	0	0
Low (27-39)	27	28.01
Moderate (40-52)	55	57.30
High (53-65)	14	14.60

Note: Categories arbitrarily made by the researcher (Cutoff scoring was determined through averaging from perfect score into Quintiles); Perfect score is 65.

Diener et al. (1985). The survey took about 10 to 15 minutes to complete. The survey was conducted where older adults congregated in groups or individually at their homes or work area.

Qualified research assistants were employed to help in the qualitative data collection. They were at least high school graduates and able to communicate and document written information from the research participants. They were initially required to undergo a training that was personally conducted by the researcher herself before the data collection process. The research assistants were also asked to sign a confidentiality agreement form.

The administration of the research questionnaires was carried out by the researcher herself and the investigator's research assistants. The researcher hired three research assistants who helped in the data gathering process. The data collection was completed

within four weeks. The participants' concerns and questions were addressed during the data collection process. The participants were also asked to indicate their agreement to participate in the research study by asking them to sign a consent form before giving them the questionnaire or allowing them to start answering the questions. Only those who were keen and willing to be included in the study were involved.

The study respondents were invited voluntarily by giving them a letter of invitation indicating the aims and purpose of the study. They were also informed of their right to refuse and it was highlighted to them that participation was voluntary. The participants were asked to sign an informed consent indicating their voluntary participation.

Data Analysis

The data gathered were tabulated and transferred to the Statistical Package for the Social Sciences for analysis (file version 1.0.0-51). Data analysis of the sample characteristics was utilized as descriptive statistics to determine the sample characteristics. The correlation of the two main variables (self-control and life satisfaction) was determined using Pearson's correlation coefficient. Correlation coefficients vary from -1 to +1, where 0 indicates no relationship, -1 indicates a perfectly negative linear relationship, and +1 indicates a perfectly positive linear relationship. For a positive coefficient, as one variable increases, the other also increases. For negative coefficients, as one variable increases, the other variable decreases (Pallant, 2020). Cohen's standard (Cohen, 2013) was used to evaluate the coefficient to assess the strength of the relationship between self-control and life satisfaction. Coefficients between 0.10 and 0.29 represent a small relationship; coefficients between 0.30 and 0.49 represent a medium relationship and coefficients 0.50 and above represent a large relationship. An alpha of 0.05 was used for analysis.

Ethical Consideration

Data collection started after approval from the Cebu Normal University-Research Ethics Committee (CNU-REC) with CNU REC Code 335/2019-06 Caorong. The study participants were purposively selected to join the study. They were screened following certain inclusion criteria such as the following: older adults aged sixty years old and above, cognitively, physically and psychologically well, and female or male residents of Iligan City and Lanao del Norte. Each of the participants was given a copy of the informed consent duly approved by the Cebu Normal University-Ethics Review Committee (CNU-REC).

Table 3 Descriptive statistics for the older adults' life satisfaction (n = 96)

Indicators	Mean	Standard Deviation	Descriptive Rating
I am satisfied with my life	5.50	1.472	Satisfied
The conditions of my life are excellent	5.44	1.609	Satisfied
In most ways my life is close to my ideal	5.34	1.368	Satisfied
So far I have gotten the important things I want in life	5.15	1.402	Slightly Satisfied
If I could live my life over, I would change almost nothing	5.07	1.643	Slightly Satisfied
Overall Mean	5.30		Satisfied

Scaling:

1.00-1.85 = Strongly Disagree/ Extremely Dissatisfied;

1.86-2.71 = Disagree/ Dissatisfied;

2.72-3.57 = Slightly Disagree;

3.58-4.43 = Neither Agree nor Disagree/ Neutral;

4.44-5.29 = Slightly Agree/ Slightly Satisfied;

5.30-6.15 = Agree/ Satisfied;

6.16-7.00 = Strongly Agree/ Extremely Satisfied

Categorization of scaling was determined through averaging from perfect score resulting to seven classified ranges.

*Perfect score is 7 in each item

Table 4 Correlation between the older adults' self-control and their life satisfaction

		Life-satisfaction
Self-control	Pearson Correlation	0.248*
	Sig. (2-tailed)	0.015
	N	96

* Correlation is significant at the 0.05 level (2-tailed).

Results

This section of the study presents the results and findings of this current study. To test the hypothesis that older adults' self-control is associated with life satisfaction, correlation statistics was performed. The presentation starts with descriptive statistics describing the characteristics of the respondents in terms of age, gender, religion, educational attainment, income and occupation. The older adults' self-control scores and their level of life satisfaction are also presented in this section of the study as well as the correlation between the two.

Table 1 indicates that the majority of the respondents who participated in the current study are young older adults, most are female Roman Catholics, and the majority of them still remain married in late life. Additionally, a great number of study participants did not reach college and more than half of them were unemployed. It can also be noted from the findings that the majority of the older participants have a monthly income of less than five thousand pesos.

Results as shown in Table 2 revealed that, generally, the respondents have moderate self-control (M=44.10, SD= 6.72) using the Brief Self-control Scale. This result is shown in the above figure, indicating that more than half of the older people involved in the study have moderate self-control. Yet it cannot be discounted from the figure above that there are fairly a number of study participants who have low level of self-control while participants of the study with high self-control are lesser in number.

It is established from the results above that the older adult participants are satisfied with life. The data show that, generally, the participants agree to the first three

indicators listed in the scale while slightly agreeing to the other two indicators.

Table 4 shows the correlation between the two main variables of the current study, which are self-control and life satisfaction. Using Pearson correlation statistics, the computed r value signifies that there is a positive relationship between the older adults' self-control and their life satisfaction. It means that a significant positive relationship between the identified variables exist. This suggests that the higher the self-control, the more satisfied the older adult is. Although Pearson's correlation between the self-control of the older adult respondents and their life satisfaction is significant at the 0.5 significance level, the magnitude is relatively small. This may be due to the sample size (N=96), hence an indication that further study must involve a much larger sample.

Discussions

The aim of this current study was to determine the link between self-control and life satisfaction among the older people to validate a proposition of the Theory of Self-control in Old age which states that the exercise of self-control leads to life satisfaction. In this study, it was hypothesized that older adults who have high self-control have better life satisfaction.

Findings of this present study reveal that the majority of the older adult participants have moderate level of self-control while are also satisfied with life. The result of the study coincides with several modern empirical studies indicating the relevance of self-control in achieving important life outcomes. Good self-control as being able to give up immediate desires and focus on long-term

goals, positively predicts life satisfaction as was concluded by Hofmann et al. (2014) in their study.

The present findings reveal that the majority of the older adult participants of the study are satisfied with life. This result is arguably plausible as was concluded by a study of Mercier, Peladeau and Tempier (1998) to older people determining their satisfaction with life. It was found in their study that, with aging, older people's psychological state remains stable or even increases even with some inevitable aging changes being experienced. Moreover, the satisfaction with life does not necessarily change in late life, as was established by Carstensen, Isaacowitz and Charles (1999). The result of this study is also in consonance with the findings of a recent study conducted by Ng, Tey and Asadullah (2017) among China's eldest people where the majority of the older adults who participated in their study considered their satisfaction with life as very good.

Furthermore, as exercising self-control has been associated with many positive outcomes, including life satisfaction, the result of the current study positively linking self-control to life satisfaction is not at all unforeseen. This is corroborated by numerous studies claiming that high self-control positively predicts not only a person's life satisfaction but also their well-being (Hofmann et al., 2014). Moreover, in a study conducted by Dou et al. (2016) to Chinese employees, self-control was also found to be positively related to both job and life satisfaction. This provides evidence of the positive association between self-control and life satisfaction. Additionally, it was also concluded in a study that high trait self-control positively predicts happiness (Cheung et al., 2014). This current study has a similar finding, that there is an apparent positive association of self-control with life satisfaction. The higher self-control the more satisfied are the older adults.

Understanding the link between self-control and life satisfaction has potential applications across a broad range of human behaviors. Linking self-control exercise to life satisfaction offers an innovative way to make people know why some may feel satisfied and why some do not.

Conclusions

This current study has provided evidence to support the proposition claiming that exercising self-control leads to life satisfaction. The significant positive relationship between the older adults' self-control exercise and their life satisfaction means that effective exercise of self-control positively leads to better life outcomes such as satisfaction in life. Linking self-control exercise to life satisfaction offers an innovative way for people to know why some people are satisfied with life and others are not.

Findings of this research underscore the positive relationship of self-control and life satisfaction among

older adults. It is recommended that further research be conducted using other variables such as quality of life to enhance the exploration concerning the relationship of self-control to the subjective well-being of the older people.

References

- Boals, A., Vandellen, M. R. and Banks, J. B. (2011) 'The relationship between self-control and health: The mediating effect of avoidant coping', *Psychology & health*. Taylor & Francis, 26(8), pp. 1049–1062.
- Bogg, T. and Roberts, B. W. (2004) 'Conscientiousness and health-related behaviors: a meta-analysis of the leading behavioral contributors to mortality', *Psychological bulletin*. American Psychological Association, 130(6), p. 887.
- Carstensen, L. L., Isaacowitz, D. M. and Charles, S. T. (1999) 'Taking time seriously: a theory of socioemotional selectivity', *American psychologist*. American Psychological Association, 54(3), p. 165.
- Cheung, T. T. L. et al. (2014) 'Why are people with high self-control happier? The effect of trait self-control on happiness as mediated by regulatory focus', *Frontiers in psychology*. Frontiers, 5, p. 722.
- Cohen, J. (2013) *Statistical power analysis for the behavioral sciences*. Routledge.
- Deary, I. J. et al. (2009) 'Age-associated cognitive decline', *British medical journal*. Oxford University Press, 339(7728), pp. 135–152.
- De Ridder, D. T. D. et al. (2012) 'Taking stock of self-control: A meta-analysis of how trait self-control relates to a wide range of behaviors', *Personality and Social Psychology Review*. Sage Publications Sage CA: Los Angeles, CA, 16(1), pp. 76–99.
- Diener, E. et al. (1999) 'Subjective well-being: Three decades of progress', *Psychological bulletin*. American Psychological Association, 125(2), p. 276.
- Diener, E. D. et al. (1985) 'The satisfaction with life scale', *Journal of personality assessment*. Taylor & Francis, 49(1), pp. 71–75.
- Dou, K. et al. (2016) 'The relationship between self-control, job satisfaction and life satisfaction in Chinese employees: A preliminary study', *Work*. IOS Press, 55(4), pp. 797–803.
- Finkel, E. J. and Campbell, W. K. (2001) 'Self-control and accommodation in close relationships: an interdependence analysis', *Journal of personality and social psychology*. American Psychological Association, 81(2), p. 263.
- Friedman, H. S. et al. (1993) 'Does childhood personality predict longevity?', *Journal of personality and social psychology*. American Psychological Association, 65(1), p. 176.
- Gilman, R. (2009) 'Shannon M. Suldo, E. Scott Huebner, Allison A. Friedrich', *Handbook of Positive Psychology in Schools*. Routledge, p. 27.
- Goodwin, R. D. and Friedman, H. S. (2006) 'Health status and the five-factor personality traits in a nationally representative sample', *Journal of health psychology*. Sage Publications Sage CA: Thousand Oaks, CA, 11(5), pp. 643–654.
- Hofmann, W. et al. (2014) 'Yes, but are they happy? Effects of trait self-control on affective well-being and life satisfaction', *Journal of personality*. Wiley Online Library, 82(4), pp. 265–277.
- Jensen-Campbell, L. A. and Malcolm, K. T. (2007) 'The importance of conscientiousness in adolescent interpersonal relationships', *Personality and Social Psychology Bulletin*. Sage Publications Sage CA: Thousand Oaks, CA, 33(3), pp. 368–383.
- Laarni, C. (2019) *Theory of Self-control in old-age*. Cebu Normal University, Philippines.
- Mercier, C., Peladeau, N. and Tempier, R. (1998) 'Age, gender and quality of life', *Community mental health journal*. Springer, 34(5), pp. 487–500.
- Ng, S. T., Tey, N. P. and Asadullah, M. N. (2017) 'What matters for life satisfaction among the oldest-old? Evidence from China', *PloS one*. Public Library of Science San Francisco, CA USA, 12(2), p. e0171799.
- Pallant, J. (2020) *SPSS survival manual: A step by step guide to data analysis using IBM SPSS*. Routledge.
- Pfeiffer, E. (1975) 'A short portable mental status questionnaire for the assessment of organic brain deficit in elderly patients', *Journal of the American Geriatrics Society*. Wiley Online Library, 23(10), pp. 433–441.
- Sayette, M. A. and Griffin, K. M. (2004) 'Self-regulatory failure and addiction', *Handbook of self-regulation: Research, theory, and applications*, 3, pp. 571–590.
- Schmeichel, B. J., Vohs, K. D. and Baumeister, R. F. (2018) 'Intellectual performance and ego depletion: Role of the self in logical reasoning and other information processing', in *Self-Regulation and Self-Control*. Routledge, pp. 310–339.

Caorong (2022)

Tangney, J. P., Boone, A. L. and Baumeister, R. F. (2018) 'High self-control predicts good adjustment, less pathology, better grades, and interpersonal success', in *Self-regulation and self-control*. Routledge, pp. 173–212.

How to cite this article: Caorong, L. A. (2022) 'The link between self-control and satisfaction in life among older adults', *Jurnal Ners*, 17(1), pp. 19–24. doi: <http://dx.doi.org/10.20473/jn.v17i1.34150>

Anemia prevention behavior in female adolescents and related factors based on Theory of Planned Behavior: A cross-sectional study

Halfie Zaqiyah Gusti Puspitasari ¹ , Ni Ketut Alit Armini ^{2,*} , Retnayu Pradanie ² , Mira Triharini ³ 

¹ Professional Nursing Program, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

² Basic Nursing Department, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

³ Advanced Nursing Department, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia

*Correspondence: Ni Ketut Alit Armini. Address: UNAIR Campus C, Mulyorejo, Surabaya, East Java, 60115, Indonesia, Email: nk.alita@fkip.unair.ac.id

Responsible Editor: Praba Diyan Rachmawati

Received: 6 July 2021 ◦ Revised: 15 April 2022 ◦ Accepted: 15 April 2022

ABSTRACT

Introduction: Female adolescent is a group prone to iron deficiency anemia due to various factors such as having monthly menstruation and poor diet. Therefore, anemia prevention behavior needs to be done from an early age to avoid prolonged health problems. The purpose of this study was to analyze the factors that correlated with anemia prevention behavior in female adolescents based on the Theory of Planned Behavior.

Methods: This study used a descriptive correlational design with a cross-sectional approach. The population of this study is female adolescents in Madiun City. Purposive sampling was used to obtain 105 respondents. The instrument used was a questionnaire to measure the variables attitude, subjective norm, perceived behavioral control, intention, and anemia prevention behavior. Data analysis was performed using the Spearman Rho Test with a level of significance ≤ 0.05 .

Results: There is a relationship between attitudes ($p = 0.003$; $r = 0.292$), subjective norms ($p = 0.006$; $r = 0.266$), and perceived behavioral control ($p = 0.002$; $r = 0.299$) with the intention to prevent anemia and also perceived behavioral control ($p = 0.003$; $r = 0.292$) and intention ($p = 0.000$; $r = 0.392$) with anemia prevention behavior.

Conclusions: Better attitude, subjective norm, perceived behavioral control, and intention possessed by a female adolescent will make better anemia prevention behavior too. Therefore, support from various parties is needed to intensify the promotion of anemia prevention, education on early detection of anemia, especially for female adolescents, and further research to determine effective health promotion methods.

Keywords: adolescent; anemia; behavior; healthcare; prevention; healthy lifestyle

Introduction

Female adolescent is a group that is prone to anemia (Andriastuti et al., 2020). Anemia is a condition of changing morphology and reducing the number of blood cells and hemoglobin so that they are not sufficient to meet the physiological needs of the body (Levy, De la Cruz Góngora and Villalpando, 2016). In female adolescents, iron deficiency anemia is the largest cause of morbidity and mortality (WHO, 2017). Several factors such as poor nutritional status, low socioeconomic status, comorbidities (malaria, thalassemia, etc.), impaired nutrient absorption, and irregular menstrual patterns will increase the chance of anemia (Abioye and Fawzi,

2020). In general, the prevalence of anemia in the world reaches 50-80% of the total population (Kassebaum et al., 2016). In Indonesia, the incidence of anemia reaches 23.7% with the prevalence of incidence in female adolescents aged 15-24 years old 32% (Ministry of Health Indonesia, 2019).

The incidence of anemia in female adolescents increases because every month they undergo menstruation, which causes an increase in iron expenditure (Swati and Sunita, 2021). In addition, many female adolescents are restricting food intake, causing an iron deficiency, and disrupting hormonal balance (Abioye and Fawzi, 2020). This is further exacerbated by the



perception of barriers such as feeling that iron tablets consumption does not provide benefits, unattractive forms of packaging and iron tablets, peers who do not consume iron tablets and dislike for the taste of iron tablets (Aprianti, Sari and Kusumaningrum, 2018).

Unsupportive behavior of female adolescents in anemia prevention efforts is influenced by knowledge levels, attitudes, perceptions, supporting environments, resources, and national policies (Roche et al., 2018). The better the support provided by the family, the more adaptive the health behaviors performed by adolescents (Annisa, Mulyono and Widyatuti, 2021). In addition, the presence of training, monitoring, availability of iron tablets, and strengthening health promotion can improve anemia prevention behavior (Gosdin et al., 2020). Some of the effects of anemia include decreased endurance, cognitive-developmental disorders, psychomotor disorders, and increased susceptibility to infection (Spezia et al., 2018). In female adolescents, the long-term impact will be more dangerous because it increases the risk of premature infant birth, bleeding, and maternal mortality (Levy, De la Cruz Góngora and Villalpando, 2016; Zekiye and Rukiye, 2021).

Based on basic health research (RISKESDAS) 2018, the prevalence of anemia in adolescents is 32%, which means that 3-4 teenagers out of 10 suffer from anemia (Ministry of Health Indonesia, 2018). According to findings, 23% of young girls and 12% of boys in East Java had anemia (Report of Office of Women's Empowerment, 2019). Through circular letter number HK.03.03/V/0595/2016 concerning the Provision of Blood Iron Tablets to Female Adolescents and Women of Childbearing Age, the government implements a program of giving one tablet per week throughout the year to female adolescents aged 12-18 years old (Ministry of Health Indonesia, 2016). In East Java Province, as many as 29.7% of female adolescents aged 10-19 years old have received iron tablets, but their consumption success rate is still low (Ministry of Health Indonesia, 2019). Handling anemia in adolescents is increasingly difficult because of behaviors such as consuming tea after eating, improper eating patterns, and consuming less iron from animal sources, causing low body mass (Mistry et al., 2017).

Implementing anemia prevention behavior from an early age is very important, especially for female adolescents to be able to prepare themselves well in facing pregnancy and childbirth. In the Theory of Planned Behavior (TPB), it is explained that the good and bad attitudes, subjective norms, and perceived behavioral control possessed by individuals will affect the amount of intention they have, resulting in differences in behavior displayed (Ajzen, 2005). The purpose of this study was to analyze factors related to anemia prevention behavior in adolescent girls based on TPB.

Materials and Methods

Study Design

This study used a descriptive correlational design through cross-sectional approach conducted in Madiun, East Java in February 2021.

Respondent

The participants involved was female adolescents selected by purposive sampling techniques. A total of 105 respondents met the inclusion criteria, and all were included in this research. The study's inclusion criteria were female adolescents aged 18-22 years old, have menstruation, and can use Google Form. The exclusion criterion is married female adolescents.

Instrument

The instrument in this study used a questionnaire prepared by the researcher, based on the TPB component and parameter, to obtain demographic data information and measure the independent variables (attitudes, subjective norms, perceptions of behavioral control, intentions) and dependent variable (anemia prevention behavior) in female adolescents. Demographic data consisted of name (initials), age, last education, age at first menstruation, menstrual pattern, knowledge about anemia, ever or not experiencing anemia, and sources of information about anemia. The number of statements on the questionnaire in each variable was attitude (n = 8), subjective norm (n = 7), perceived behavioral control (n = 7), intention (n = 6), and anemia prevention behavior (n = 7). All questions were measured using a 5-point Likert scale with answer choices on the independent variable (1 = strongly disagree; 2 = disagree; 3 = undecided; 4 = agree; 5 = strongly agree) and the dependent variable (1 = never; 2 = rarely; 3 = sometimes; 4 = often; 5 = always). The final results on the scoring were categorized into good, enough, and less. The instrument has been tested on 20 young women with an age range of 18-24 to get the results of the validity test ($r > 0.444$) and reliability (Cronbach's alpha attitude = 0.733; subjective norm = 0.704; perceived behavioral control = 0.636; intention = 0.727, anemia prevention behavior = 0.617).

Data Collection

Data collection used online questionnaire by compiling a list of prospective respondents based on criteria and large samples that have been determined from female adolescents who are members of the Madiun student organization. Furthermore, the researchers explained the research to respondents and examined the Google Form filled out by the respondent to ensure that the data submitted were appropriate and recorded in the researcher's database.

Data Analysis

The data were analyzed using the descriptive analysis method to determine the percentage and frequency distribution of the data and bivariate analysis to

Table 1 Demographic characteristics of respondents (N = 105)

Characteristics	Parameter	n	%	Mean	SD
Age	18 years old	7	6.7	20.11	1.13
	19 years old	28	26.7		
	20 years old	29	27.6		
	21 years old	28	26.7		
	22 years old	13	12.4		
Last education	Senior High School	98	93.3	12.32	1.28
	Diploma	5	4.8		
	Bachelor	2	1.9		
Age at first menstruation	≤12 years old	62	59.1	12.32	1.28
	>12 years old	43	40.9		
Length of menstruation	3-4 days	2	1.9	NA	NA
	4-5 days	21	20		
	5-6 days	22	21		
	6-7 days	50	47.6		
	>7 days	10	9.5		
Many change of sanitary pad in a day	2 times	13	12.4	3 times	0.88
	3 times	53	50.5		
	4 times	24	22.9		
	>4 times	15	14.3		
Know anemia	Yes	103	98.1	NA	NA
	No	2	1.9		
Have anemia	No	53	50.5	NA	NA
	Yes	52	49.5		
Source of anemia information	Family	17	16.2	NA	NA
	University	12	11.4		
	Social Media	25	23.8		
	School	47	44.8		
	Book	1	1		
	Internet	2	1.9		
Know the guidelines for prevention and control of anemia	Yes	10	9.5	NA	NA
	No	95	90.5		

Table 2 Relationship between attitudes, subjective norms, and perceptions of behavioral control with the intention to prevent anemia in female adolescent (N = 105)

Variable	Category	Intention						Spearman Rho Test	
		Less		Enough		Good		p	r
		n	%	n	%	n	%		
Attitude	Positive	7	6.7	28	26.7	18	17.1	0.003	0.292
	Negative	12	11.4	36	34.3	4	3.8		
Subjective Norm	Good	3	2.9	10	9.5	12	11.4	0.006	0.266
	Enough	12	11.4	42	40.0	8	7.6		
	Less	4	3.8	12	11.4	2	1.9		
Perceived Control Behavior	Good	2	1.9	17	16.2	14	13.3	0.002	0.299
	Enough	17	16.2	40	38.1	7	6.7		
	Less	0	0	7	6.7	1	1.0		

Table 3 Relationship between perceived behavioral control and intention with anemia prevention behavior in female adolescent (N=105)

Variable	Category	Anemia Prevention Behavior						Spearman Rho Test	
		Less		Enough		Good		p	r
		n	%	n	%	n	%		
Perceived Control Behavior	Good	4	3.8	24	22.9	5	3.8	0.003	0.292
	Enough	23	21.9	34	32.4	7	6.7		
	Less	5	4.8	3	2.9	0	0		
Intention	Good	2	1.9	16	15.2	4	3.8	<0.001	0.392
	Enough	17	16.2	39	37.1	8	7.6		
	Less	13	12.4	6	5.7	0	0		

determine the relationship between the independent and dependent variables using the Spearman Rho Test with $\alpha \leq 0.05$. If the value of $p < 0.05$ then H_0 was rejected. It means there is a relationship between attitude, subjective norm, and perceived control behavior with the intention to prevent anemia, also there is a relationship between perceived control behavior and intention with anemia prevention behavior.

Ethical Consideration

This research has been reviewed and approved by the Health Research Ethics Commission, Faculty of Nursing, Universitas Airlangga with No. 2166-KEPK. The ethical components considered in this study are informed consent, anonymity, and confidentiality.

Results

The majority of respondents in this study were 20 years old (27.6%) and the last education level was high school, namely 98 respondents (93.3%). In addition, as many as 62 respondents (59.1%) experienced their first menstruation at the age of 12 years old. Most of the young women in this study experienced menstruation for 6-7 days each month (47.6%). During menstruation they changed sanitary pad twice (12.4%), three times (50.5%), four times (22.9%), and more than four times (14.3%). Regarding knowledge about anemia, 103 respondents (98.1%) knew about anemia and 53 respondents (50.5%) said that they had anemia. Schools are the main source of information for a female adolescent to gain knowledge about anemia, as many as 47 respondents (44.8%), followed by social media ($n = 25$; 27.8%), and families ($n = 17$; 16.2%). In addition, 95 respondents (97.5%) said they never knew about the Guidelines for Prevention and Management of Anemia, a guide in the form of a module issued by the Ministry of Health for adolescents and women of childbearing age to deal with anemia (Table 1).

Based on Table 2, it can be seen that there is a relationship between attitudes ($p = 0.003$; $r = 0.292$), subjective norms ($p = 0.006$; $r = 0.266$), and perceived behavioral control ($p = 0.002$; $r = 0.299$) with intention to prevent anemia and the relationship created has a moderate degree of strength. In the attitude variable, the highest number is found in respondents with negative attitudes and having enough intention to prevent anemia, namely 36 respondents (34.3%). Furthermore, in the subjective norm variable, the highest number is found in respondents with enough subjective norms and having enough intention to prevent anemia, as many as 42 respondents (40%). Meanwhile, in the perceived behavioral control variable, the highest number was found in respondents with enough behavioral control perceptions and had sufficient intention to prevent anemia, namely 40 respondents (38.1%).

Based on Table 3, it can be seen that there is a relationship between perceptions of behavioral control ($p = 0.003$; $r = 0.292$) and intentions ($p < 0.001$; $r = 0.392$) with anemia prevention behavior and the relationship created has a moderate degree of strength. In the behavioral control perception variable, the highest number is found in respondents with enough behavioral control perceptions and realizing sufficient anemia prevention behavior as many as 34 respondents (32.4%). Furthermore, in the intention variable, the highest number is found in respondents with enough intention or intention and realizing sufficient anemia prevention behavior, as many as 39 respondents (37.1%).

Discussions

Most female adolescents have enough behavior in preventing anemia and some are already in the good category. However, it was still found that some

respondents fall into the category of lacking in anemia prevention behavior. Behavioral differences embodied by these female adolescents can be influenced by several factors, namely, age, gender, health information received, knowledge, and income of parents to affect the level of health knowledge due to low interest in health literacy (Fleary, Joseph and Pappagianopoulos, 2018). Behavior is also influenced by repeated habits that form a cycle (Jalambo et al., 2018). A person will display adaptive behavior when they know the benefits, have sufficient pressure or social drive, and support, both internal and external. TPB relates to attitudes, subjective norms, and perceived behavioral control that affect the magnitude of the intention to realize health behavior (Ajzen, 2005). The main findings of this research show that attitudes, subjective norms, perceived behavioral control, and intention have strong relationship with anemia prevention behavior among female adolescents.

There is a significant relationship between the attitude of female adolescents to prevent anemia. Adolescents with a positive attitude will have better abilities in realizing health (Guerin and Toland, 2020). However, not all respondents with a positive attitude have good intentions in preventing anemia. Some of them still have a negative attitude but have enough and good intentions to prevent anemia. This is related to the low level of knowledge of young women about signs and symptoms and how to prevent anemia. Someone who knows is not necessarily able to realize the information they get in the form of behavior (Fleary, Joseph and Pappagianopoulos, 2018). Another factor that causes this to happen is the age of the respondent in the adolescent stage, making decisions about an action to be taken is influenced by differences in the level of psychological maturity, cognitive abilities, and symptoms of depression (McCue et al., 2019). Health workers are an important part of being able to increase awareness and modify interventions for female adolescent so that they have knowledge, attitudes, and behaviors that support preventing anemia (Nesrin, Anwar and Abdullah, 2021).

Subjective norms are also related to the intention to prevent anemia. This result is in line with research that states that adolescents have the intention to manifest health behaviors when they have high subjective norm support, especially from parents and peers (Zhao et al., 2020). The incidence of anemia will increase two times higher if adolescents only live with one of their parents (father or mother) and 2.4 times higher if they live with their guardian (Gonete et al., 2018). Adolescents who receive emotional and instrumental support from teachers and friends will also be more concerned with their health condition because they have a positive influence (Triana et al., 2019). The attention given by the people closest to them or those who are considered important and authorized, may not necessarily change the perspective of female adolescents to influence the intention to prevent anemia. In addition, adolescents

have a high sense of ego so that they seem unruly and act selfishly, and do not think about the impact of the choices they make (Krisnani and Farakhiyah, 2017).

On the other hand, there is a relationship between perceived behavioral control and the intention to prevent anemia. Adolescents with a strong perception of behavioral control will have 3.906 times higher intention to comply with iron tablet supplement consumption (Quraini, Ningtyias and Rohmawati, 2020). Beliefs in the perception of behavioral control can be influenced by the support received and the habits that are carried out to form a recurrent value (Triana et al., 2019). Most of the respondents in this study have a perception of sufficient behavioral control and sufficient intensity of anemia prevention. This is due to the ability to meet the body's lack of nutrition to prevent anemia, especially during menstruation. In fact, during menstruation, adolescent girls will lose hemoglobin more quickly, so they need to increase their intake of nutrients, especially iron (Masoud et al., 2020). However, this will be difficult if young women have low socioeconomic conditions, because they are related to the fulfillment of nutrition from the food that can be consumed daily (Banayjeddi et al., 2019).

Perceived behavioral control is also directly related to anemia prevention behavior. Some of the female adolescents in this study already had a good level of perception of behavioral control so that they would carry out anemia prevention behavior well. However, some female adolescents have a perception of sufficient behavioral control but can perform anemia prevention behavior well or vice versa. This is influenced by family support, information, communication, and the level of emotional quality (Park and Lee, 2020). In addition, the emergence of opportunities for adaptive behavior is higher when adequate health facilities are available. If individuals are aware that the perceived ease of accessing health services is higher than the barriers, it will form a positive perception of being able and utilizing the available facilities in realizing health behaviors (Roche et al., 2018).

Intention also has a significant relationship with anemia prevention behavior. In TPB, there is a base rate term which states that individuals will tend to perform behaviors that are mostly done by others, and vice versa (Ajzen, 2005). In addition, the good or bad health behavior that will be realized is also strongly influenced by various kinds of life contexts such as the environment, skills, and existing health programs (Nagy-Pénczes, Vincze and Biro, 2020). Good intentions can arise because of health-related actions taken every day. Female adolescents who regularly consume vegetables and fruit every day will meet their vitamin and iron needs (Ahankari et al., 2017). Awareness of the consumption of vegetables and fruit will arise from the knowledge that the older you get, the more your iron needs will increase (Masoud et al., 2020). Wrong dietary habits and consumption of unhealthy food (junk food) should be

avoided (Jalambo et al., 2018). Female adolescents must also meet the necessary nutrients, not only macro but also micro (Htet et al., 2016). In addition, female adolescents who get iron tablets from school or have independent supplies tend to consume iron tablets regularly (Gosdin et al., 2020). This habit will directly or indirectly foster intention and shape anemia prevention behavior.

The limitation of this research is that it is a cross-sectional study that only emphasizes the measurement of variables at one time so that there is no follow-up on the results of observations. The parameters used in the study are few so that they have not been able to provide a more detailed description of the respondent's condition. In addition, the online data collection process made the researcher unable to see the respondent's physical condition (height, weight, signs of anemia, etc.), if they did not read the explanation, as well as the instructions for filling out the questionnaire properly.

Conclusions

The better intention possessed by adolescent girls, the better anemia prevention behavior is carried out, while attitudes, subjective norms, and perceived behavioral control are factors that affect the quality of the intentions. The factor that most strongly influenced the quality of intention was the perceived behavioral control, while the most influential support came from the family. It is necessary to increase knowledge about anemia prevention behavior to female adolescents and their families as the main source of support for adolescents in adaptive behavior. In addition, further research is expected to be able to find effective methods in increasing knowledge and awareness for young women in realizing anemia prevention behavior from an early age.

References

- Abioye, A. I. and Fawzi, W. W. (2020) 'Nutritional Anemias', *Present Knowledge in Nutrition (Eleventh Edition)*. Boston, 2, pp. 503–521. doi: 10.1016/B978-0-12-818460-8.00027-7.
- Ahankari, A. S. et al. (2017) 'Prevalence of Iron-deficiency Anaemia and Risk Factors in 1010 Adolescent Girls from Rural Maharashtra, India: A Cross-Sectional Survey', *Public Health*. Elsevier Ltd, 142, pp. 159–166. doi: 10.1016/j.puhe.2016.07.010.
- Ajzen, I. (2005) *Attitudes, Personality, and Behavior*. Second Edi. United Kingdom: Open University Press.
- Andriastuti, M. et al. (2020) 'Prevalence of anemia and iron profile among children and adolescent with low socio-economic status', *International Journal of Pediatrics and Adolescent Medicine*. Elsevier Ltd, 7(2), pp. 88–92. doi: 10.1016/j.ijpam.2019.11.001.
- Annisa, F., Mulyono, S. and Widayati, W. (2021) 'Family Support on Utilization of Adolescent Reproduction Health Services at The Area of Public Health Service (Puskemas) of Martapura', *Enfermeria Clinica*, 31(2), pp. S153–S138. doi: doi.org/10.1016/j.enfcli.2020.12.008.
- Aprianti, R., Sari, G. M. and Kusumaningrum, T. (2018) 'Factors Correlated with the Intention of Iron Tablet Consumption among Female Adolescents', *Jurnal Ners*, 13(1), p. 122. doi: 10.20473/jn.v13i1.8368.
- Banayjeddi, M. et al. (2019) 'Implementation evaluation of an iron supplementation programme in high-school students: The

- crosswise model', *Public Health Nutrition*, 22(14), pp. 2635–2642. doi: 10.1017/S1368980019001575.
- Fleary, S. A., Joseph, P. and Pappagianopoulos, J. E. (2018) 'Adolescent Health Literacy and Health Behaviors: A Systematic Review', *Journal of Adolescence*. Elsevier, 62(March 2017), pp. 116–127. doi: 10.1016/j.adolescence.2017.11.010.
- Gonete, K. A. et al. (2018) 'Prevalence and Associated Factors of Anemia among Adolescent Girls Attending High Schools in Dembia District, Northwest Ethiopia, 2017', *Archives of Public Health*. Archives of Public Health, 76(1), pp. 1–9. doi: 10.1186/s13690-018-0324-y.
- Gosdin, L. et al. (2020) 'Barriers to and Facilitators of Iron and Folic Acid Supplementation within a School-based Integrated Nutrition and Health Promotion Program among Ghanaian Adolescent Girls', *Current Developments in Nutrition*. Oxford University Press, 4(9), pp. 1–11. doi: 10.1093/cdn/nzaa135.
- Guerin, R. J. and Toland, M. D. (2020) 'An Application of A Modified Theory of Planned Behavior Model to Investigate Adolescents' Job Safety Knowledge, Norms, Attitude and Intention to Enact Workplace Safety and Health Skills', *Journal of Safety Research*. National Safety Council, 72(December), pp. 189–198. doi: 10.1016/j.jsr.2019.12.002.
- Htet, M. K. et al. (2016) 'Folate and Vitamin B 12 Status and Dietary Intake of Anaemic Adolescent Schoolgirls in the Delta Region of Myanmar', 116, pp. 36–41. doi: 10.1017/S0007114515001609.
- Jalambo, M. O. et al. (2018) 'Prevalence and Risk Factor Analysis of Iron Deficiency and Iron-deficiency Anaemia among Female Adolescents in the Gaza Strip, Palestine', (8), pp. 1–10. doi: 10.1017/S1368980018001568.
- Kassebaum, N. J. et al. (2016) 'The Global Burden of Anemia', *Hematology/Oncology Clinics of North America*. Elsevier Inc, 30(2), pp. 247–308. doi: 10.1016/j.hoc.2015.11.002.
- Krisnani, H. and Farakhiyah, R. (2017) 'Meningkatkan Kemampuan Pengambilan Keputusan pada Remaja Akhir dengan Menggunakan Metode Realty Therapy', *Share Social Work Jurnal*, 7(2), pp. 1–79.
- Levy, T. S., De la Cruz Góngora, V. and Villalpando, S. (2016) 'Anemia: Causes and Prevalence', *Encyclopedia of Food and Health*, pp. 156–163. doi: 10.1016/B978-0-12-384947-2.00029-5.
- Masoud, M. S. et al. (2020) 'The association between iron and Vitamin D status in Arab adolescents', *Public Health Nutrition*, 23(7), pp. 1208–1213. doi: 10.1017/S1368980019001113.
- Mccue, R. et al. (2019) 'The Future and Me: Imagining the Future and the Future Self in Adolescent Decision Making', *Cognitive Development*. Elsevier, 50(December 2018), pp. 142–156. doi: 10.1016/j.cogdev.2019.04.001.
- Ministry of Health Indonesia (2016) *Guidelines for the Prevention and Management of Anemia in Adolescent Girls and Women of Childbearing Age (WUS)*. Directorate of Community Nutrition and Directorate General of Public Health, Ministry of Health, Republic of Indonesia.
- Ministry of Health Indonesia (2018) *Report Basic Health Research 2018*. Jakarta.
- Ministry of Health Indonesia (2019) *National Report Basic Health Research 2018*. Publishing Institute for Health Research and Development Agency.
- Mistry, S. K. et al. (2017) 'An Outline of Anemia Among Adolescent Girls in Bangladesh: Findings from a Cross-sectional Study', *BMC Hematology*. BMC Hematology, 17(1), pp. 1–8. doi: 10.1186/s12878-017-0084-x.
- Nagy-p, G., Vincze, F. and B, É. (2020) 'Does Better Health-Related Knowledge Predict Favorable Health Behavior in Adolescents?', 17.
- Nesrin, N. A.-B., Anwar, M. E. and Abdullah, M. K. (2021) 'The Impact of Nutrition Education on Knowledge, Attitudes, and Practice Regarding Iron Deficiency Anemia among Female Adolescent Student in Jordan', *Heliyon*, 7(2). doi: 10.1016/j.heliyon.2021.e06348.
- Park, H. and Lee, K. S. (2020) 'The Association of Family Structure with Health Behavior, Mental Health, and Perceived Academic Achievement among Adolescents: A 2018 Korean Nationally Representative Survey', *BMC Public Health*. BMC Public Health, 20(1), pp. 1–10. doi: 10.1186/s12889-020-08655-z.
- Quraini, D. F., Ningtyias, F. W. and Rohmawati, N. (2020) 'Compliance Behavior of Iron Tablet Supplement Consumption to Adolescent Girls in Jember, Indonesia', *The Indonesian Journal of Health Promotion and Health Education*, 8(2), pp. 154–162. doi: 10.20473/jpk.V8.I2.2020.154-162.
- Report of Office of Women's Empowerment, C. P. and P. of E. J. P. (2019) *No Tit*.
- Roche, M. L. et al. (2018) 'Adolescent Girls' Nutrition and Prevention of Anaemia: A School Based Multisectoral Collaboration in Indonesia', *BMJ (Online)*, 363, pp. 1–6. doi: 10.1136/bmj.k4541.
- Spezia, J. et al. (2018) 'Prevalence of Anemia in Schools of the Metropolitan Region of Curitiba, Brazil', *Hematology, Transfusion and Cell Therapy*. Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular, 40(2), pp. 151–155. doi: 10.1016/j.htct.2017.11.007.
- Swati, R. and Sunita, M. (2021) 'Menstrual Status and Obstetrical History Influencing Prevalence of Iron Deficiency Anemia among Reproductive Age Women in Rural Area', *Materials Today: Proceedings*. doi: https://doi.org/j.matpr.2020.12.966.
- Triana, R. et al. (2019) 'Understanding the protective factors (self-esteem, family relationships, social support) and adolescents' mental health in Jakarta', *Enfermeria Clinica*. Elsevier España, S.L.U., 29(Insc 2018), pp. 629–633. doi: 10.1016/j.enfcli.2019.04.096.
- WHO (2017) *Global Accelerated Action for the Health Adolescent: Guidance to Support Country Implementation*.
- Zekiye, K. and Rukiye, D. (2021) 'The Impact of Adolescent Pregnancy on Maternal and Infant Health in Turkey: Systematic Review and Meta-analysis', *Journal of Gynecology Obstetrics and Human Reprofusion*, 50(4). doi: 10.1016/j.jogoh.2021.102093.
- Zhao, X. et al. (2020) 'Intention to Drink and Alcohol Use Before 18 Years Among Australian Adolescents: An Extended Theory of Planned Behavior', *Addictive Behaviors*. Elsevier, 111(February), p. 106545. doi: 10.1016/j.addbeh.2020.106545.

How to cite this article: Puspitasari, H. Z. G., Armini, N. K. A., Pradanie, R., Triharini, M. (2022) 'Anemia prevention behavior in female adolescents and related factors based on Theory of Planned Behavior: A cross-sectional study', *Jurnal Ners*, 17(1), pp. 25-30. doi: <http://dx.doi.org/10.20473/jn.v17i1.27744>

Knowledge and attitudes toward COVID-19 vaccination among student nurses from Saudi Arabia

Romeo Mostoles Jr. ^{1,*} , Richard Maestrado ¹ , Joyce Buta ¹ , Hamdan Mohammad Albaqawi ¹ , Evalynne Rondilla ¹ , Salman Alsaqri ¹ , Sandro Villareal ¹ 

¹ College of Nursing, University of Hail, Hail, Saudi Arabia

*Correspondence: Romeo Mostoles Jr. Address: College of Nursing, University of Hail, Hail, Saudi Arabia, Email: r.mostoles@uoh.edu.sa

Responsible Editor: Retnayu Pradanie

Received: 10 March 2022 ◦ Revised: 12 April 2022 ◦ Accepted: 13 April 2022

ABSTRACT

Introduction: Insights into the knowledge and attitudes of students can be achieved by analyzing their general understanding and prevention of the coronavirus. Therefore, it is imperative to identify attributes that influence the development of healthy behaviors among students. This study aimed to determine the dynamics of knowledge and attitudes of student nurses concerning the COVID-19 vaccine.

Methods: This descriptive-cross sectional study involved 250 student nurses from the University of Hail, Nursing College, selected by convenience sampling. The data were collected between August and September 2021 using questionnaires. The data were analysed using a t-test, one-way ANOVA, and Pearson’s correlation test with a significance level of 0.05.

Results: The study results showed that participants have good knowledge (mean = 3.09; SD = 0.88) and attitudes (mean = 8.62; SD = 1.83). Gender (except for knowledge), marital status, and age were not relevant as explanatory factors of knowledge and attitude. There was a significant difference in both knowledge (F = 2.233; p < 0.005) and attitude (F = 3.918; p < 0.004) associated with the course year. There was no significant relationship between knowledge and attitude (r = -0.013; p-value = 0.834).

Conclusions: This information is critical for educators as they design techniques to help students put their knowledge into practice. In this way, educators may help health students develop skills and encourage students to become accountable health professionals who can support current societal healthcare demands.

Keywords: COVID-19; demographics; intention; student nurses; vaccine

Introduction

The most effective preventive measure to reduce the pandemic is predicted to be the coronavirus disease-2019 (COVID-19) vaccination (Szmyd *et al.*, 2021), and efforts have been made to produce the vaccines, especially in developing countries (Albaqawi *et al.*, 2021). Vaccination has had a 67% acceptance rate, despite the considerable demographic and geographic differences in vaccination uptake (Malik *et al.*, 2020) due to safety concerns. It is important to note that the success of a vaccination program is determined by the vaccination uptake by the public, healthcare workers (Kassianos *et al.*, 2018), and students (e.g., nursing students), who are critical advocates of vaccination. In addition, the campus

environments of student nurses are often congested with intense movement and restricted space, i.e., environments that are considered high-risk during the COVID-19 pandemic. Therefore, nursing students also benefit from vaccination. Conversely, vaccination is a critical intervention with the potential to significantly limit the occurrence and spreading of this deadly infectious illness (Jiang *et al.*, 2021).

SARSCoV-2 vaccines became available at the end of 2020, and healthcare workers (HCWs) were among the first to be vaccinated in many countries (Szmyd *et al.*, 2021). Nursing students, in turn, cover inevitable shortfalls in the healthcare sector and also need vaccination as they have played a key role in the current response to the pandemic whenever possible (Fontenot *et*



al., 2021). However, university students' desire to become immunized to COVID-19 varies greatly (Jiang *et al.*, 2021), which could be the result of a lack of awareness that leads to the non-intention of being vaccinated. Indeed, safety issues and unfortunate events have been reported throughout the implementation of vaccination programs, damaging public opinion and influencing vaccination intentions (Grady and Mazzei, 2021). The resistance to COVID-19 immunization may increase the danger of pathogen transmission to the patients of healthcare workers while simultaneously reducing the chance of encouraging patients to get vaccinated (Biswas *et al.*, 2021). Indeed, this can be due to the safety of the vaccine, its efficacy, and potential side effects. Accordingly, vaccination has been explored in the literature; however scarcely concerning the dynamics of knowledge, attitudes, and practices associated with COVID-19 among student nurses as advocate partners of the healthcare professionals.

The negative views and skepticism toward COVID-19 vaccines are key obstacles to expanding global vaccination coverage (Paul *et al.*, 2021). Nursing students are partners in supporting the vaccination programs for COVID-19. Thus, it is critical to understand the knowledge, attitudes, and practices of student nurses, helping them develop mitigation solutions for the current scenario. Most studies on the preventive behavior of nursing students during the COVID-19 pandemic have shown a high level of commitment to preventive measures (Sun *et al.*, 2020). Moreover, vaccination acceptance rates among nursing students have been reported as satisfactory (Jiang *et al.*, 2021). It is significant, therefore, that deeper insights into existing students' knowledge and attitudes be achieved by analyzing their general understanding and prevention of the coronavirus. Hence, it is imperative to assist in the identification of attributes that influence the students in the development of healthy behaviors. This study aimed to determine the dynamics of knowledge and attitudes of student nurses toward the COVID-19 vaccine. To the best of our knowledge, this is the first study that explores the knowledge and attitudes of student nurses from the Hail region, Kingdom of Saudi Arabia.

Materials and Methods

Study Design

This study used the descriptive-cross sectional approach to determine the dynamics of knowledge and attitudes among student nurses concerning the COVID-19 vaccine.

Respondent

This study involved the active participation of 250 student nurses from the University of Hail, Nursing College. A non-probability sampling, specifically convenience sampling, was used. All full-time students

were invited to participate in the study, and those who did not provide consent were excluded. The researchers used convenience sampling, which means that all students were invited to participate in the study

Instrument

The questionnaire comprised two parts. The first part consisted of the demographic characteristics, which include gender, marital status, age, and course year. The second part was adapted from Islam *et al.* (2021) and had 11 items categorized as knowledge (five items) and attitudes (six items).

The knowledge items asked about the general knowledge about the COVID-19 vaccine (e.g. "Do you know about the COVID-19 vaccine?"; "Do you know about the effectiveness of the COVID-19 vaccine?") and had three possible responses (i.e., "Yes," "No" or "Don't know"). The "Yes" response was assigned a score of 1, while the "No/Don't know" responses were scored 0. The total score ranged from 0 to 5 and was calculated by adding the raw scores of the five items, with the highest score indicating increased level of knowledge of COVID-19 vaccines.

The attitude section consisted of six items (e.g., "The newly discovered COVID-19 vaccines are wise"; "The COVID vaccines are essential for us") that can be answered through a three-point Likert scale ranging from 0 to 2, where 0 = Disagree, 1 = Undecided, and 2 = Agree. The total score was determined by adding the raw scores of the six items, which ranged from 0 to 12, with a higher overall score suggesting more favorable attitudes toward the COVID-19 vaccination.

The questionnaire was subjected to content validation by three panel experts of the Hail region, Saudi Arabia. These three experts unanimously agreed on the validity of the questionnaire. The questionnaire was subsequently tested with 15 student nurses who were no longer part of the sample. Reliability results were high (0.86 for knowledge and 0.89 for attitudes).

Table 1 Demographic characteristics of the participants (N = 250)

Demographic characteristics	n	%
Age		
20-22	118	47.2
23-25	101	40.4
26-28	20	8
29 years old and above	11	4.4
Gender		
Male	124	49.6
Female	126	50.4
Marital Status		
Single	246	98.4
Married	4	1.6
Year Level		
Level 5	50	20
Level 6	50	20
Level 7	50	20
Level 8	40	16
Internship	60	24

Table 2 Differences between demographic characteristics and knowledge and attitudes of the nursing students toward COVID-19 vaccine

Demographic characteristics	Knowledge				Attitude			
	Mean	Std. Deviation	t	P-value	Mean	Std. Deviation	t	P-value
Age								
20-22	3.00	0.901	3.673	0.073	8.93	1.884	2.623	0.071
23-25	3.06	0.863			8.33	1.866		
26-28	3.70	0.732			8.10	1.252		
29 years old and above	3.09	0.700			8.90	1.300		
Gender								
Male	2.94	0.895	-2.678	0.008*	8.77	1.928	1.287	0.199
Female	3.23	0.843			8.47	1.728		
Marital Status								
Single	3.09	0.884	0.783	0.434	8.63	1.839	0.686	0.494
Married	2.75	0.500			2.66	0.456		
Year Level								
Level 5	3.03	0.920	2.233	0.005*	7.93	1.550	3.918	0.004*
Level 6	3.00	0.890			9.33	1.660		
Level 7	3.40	0.672			9.10	1.450		
Level 8	3.09	0.641			8.90	1.300		
Internship	3.32	0.701			9.90	0.850		

* < 0.05

Data Collection

The researchers invited the students to participate through classroom announcements. The students were instructed as to what the study entails, extent of their participation, and their rights as participants, before deciding whether or not to participate in the study. Informed consent forms were distributed to the willing participants thereafter. Data were collected between August and September 2021.

Data Analysis

The SPSS v.22 was used to analyze the data. The demographic data were described using frequency and percentage values. A t-test was used to identify differences between gender and marital status for knowledge and attitude, while a one-way ANOVA was employed to determine differences between age and course year. Pearson's r was used to determine the relationship between knowledge and attitudes. The significance level of 0.05 was considered for the statistical analyses.

Ethical Consideration

The Institutional Review Board of the University of Hail has approved this protocol (H-2021-067). In the conduct of human study, the researchers followed the ethical guidelines outlined in the World Medical Association Declaration of Helsinki. All participants signed a written informed consent form.

Results

The participants were generally young, with 47.2% aged between 20 and 22 years. Gender was nearly equally distributed with males comprising 49.6% of the participants and females 50.4% (Table 1). Only 1.6% of the students were married, and most belonged to an internship program (24%).

The knowledge of the student nurses was good (3.09 out of 5), and they had a good attitude (8.62 out of 12) toward the COVID-19 vaccine (Table 2). There was a significant difference in knowledge ($t = -2.678$; $p < 0.008$) between the genders of the nursing students, such that female students had better knowledge (3.23 ± 0.843) than their male colleagues (2.94 ± 0.895). However, there were no differences in attitude between genders ($t = 1.287$; $p > 0.199$). In addition, neither knowledge ($t = 0.783$; $p > 0.434$) nor attitude ($t = 0.686$; $p > 0.494$) differed regarding the marital status of the participants. Furthermore, there was no significant difference in the knowledge ($F = 3.673$; $p > 0.703$) or attitude ($F = 2.623$; $p > 0.071$) of the nursing students associated with their age.

There was a significant difference in both knowledge ($F = 2.233$; $p < 0.005$) and attitude ($F = 3.918$; $p < 0.004$) associated with the course year (Table 2). There was no significant relationship between knowledge and attitude ($r = -0.013$; $p\text{-value} = 0.834$).

Discussions

This study aimed to determine the knowledge and attitudes of student nurses toward the COVID-19 vaccine. The knowledge of the student nurses recorded in this study was good, which can be credited to their ongoing study and their active participation in disseminating preventive information. The results of this study are similar to that of an earlier study where the mean knowledge score was 2.83 out of 5 (Islam *et al.*, 2021), also showing the good attitude of the student nurses toward COVID-19 vaccination. A positive attitude is a key to containing the outbreak and increasing willingness to receive the COVID-19 vaccine. This shows the students' willingness to be vaccinated and promote vaccination within their community or among their clients. Overall, these good results concerning the knowledge and attitudes of the student nurses show the professional foundation and basic awareness of the necessity, effectiveness, and safety of the COVID-19 vaccine. One

good example to strengthen their knowledge and attitudes is for the students to be given an opportunity to disseminate the value of having the vaccine to their clients or patients during their community and hospital practice.

There was a considerable disparity in knowledge between the genders, where females had greater scores than males. Previous research shows that females have a higher notion of self-education and self-control (Cornwell, Mustard and Van Parys, 2013), and, thus, performed better than male students. Such result is comparable to an earlier study (Pasay-An *et al.*, 2021), and shows that male students need to be more educated regarding the COVID-19 vaccine to perform their task in promoting the vaccination program. It is vital to emphasize, however, the importance of considering educational needs while developing educational initiatives, regardless of demographic considerations (Pasay-An *et al.*, 2021). This will help the student nurses better grasp their roles in promoting the COVID-19 vaccine. For example, nurse educators can encourage their student nurses to volunteer, either giving vaccine or doing health education in the vaccination area.

Marital status was not associated with the knowledge and attitudes toward the COVID-19 vaccine. Al-Hanawi *et al.* (2020) also reported that the marital status of participants was not significantly associated with knowledge and attitudes. While the result is not significant, there still needs a continuous effort to include all students (regardless of marital status) in the ongoing improvement plan for vaccine information dissemination. For example, offering and giving them an educational session and educational outreach materials to raise their public understanding on the value of vaccine to combat the disease. Meanwhile, there were also no significant differences in the knowledge and attitude among the nursing students of different ages. However, age was previously reported as an indicator of knowledge and attitudes toward the COVID-19 vaccine (Al-Hanawi *et al.*, 2020; Pasay-an, Magwilang and Pangket, 2020). A significant difference in knowledge and attitude was recorded regarding the course year, which agrees with the findings of Kumar, Pinky and Nurudden (2021) and indicates that students needed an educational program created for their specific needs.

Apparently, knowledge and attitudes are not related to one another, which agrees (Pasay-An *et al.*, 2021) and disagrees (Al-Hanawi *et al.*, 2020) with previous findings. This study found no significant connections between knowledge and attitudes concerning COVID-19, which implies that information does not necessarily convert into attitudes and/or behaviors, and that behaviors do not transcend into attitudes. One possible explanation is that student nurses have only recently learned about the vaccination and are currently making it work in their perspectives. In contrast, Erfani *et al.* (2020) discovered that increased knowledge of COVID-19 was associated

with positive attitudes and excellent practices, suggesting that good knowledge leads to positive attitudes and good practices. In context, in developing the policy, the policy makers should consider that individual knowledge and attitudes are utilized to appraise events and their potential, as well as their repercussions.

Despite the fact that this study was conducted in a local context and the results are not generalizable, the findings are nevertheless relevant in the global arena. International policy makers, for example, can use the data from this study to plan certain initiatives based on the similar features found in this study. Furthermore, the data in this study can be utilized to evaluate and compare to data from other countries in reducing the gap on this topic.

Policy implication

This research has substantial policy implications. With the findings, policy makers in a university institution can adjust the instructional intervention to the vaccine's information drive. Moreover, this study contributes to the identification of relevant communication networks for the target demographic as part of the preventative program strategy. As a result, school or university officials can focus their efforts on programs geared at minimizing the spread of misinformation and conspiracy theories about the COVID-19 vaccine. Furthermore, the university has the ability to take unprecedented efforts and respond quickly in establishing tight control and preventative measures against COVID-19 to safeguard student safety.

Strengths and weakness of the study

This study targeting the knowledge and attitudes toward COVID-19 vaccination among the student nurses provides a useful tool for gauging vaccine acceptability. In addition, it provides the policy makers with relevant recommendations. However, it has substantial limitations. For example, the study was conducted in one setting and the use of non-probability sampling means it may not be possible to generalize. Also, the non-inclusion of practice of students in the study can be more explored as the researchers failed to do so.

Conclusions

Student nurses have a good knowledge and attitude toward the COVID-19 vaccine, and female students are more knowledgeable. Knowledge and attitude did not differ significantly between gender (except for knowledge), marital status, and ages. Lastly, a significant difference in knowledge and attitude was recorded for course year. This information is critical for educators as they design techniques to help students put their

knowledge into practice. Indeed, in this way educators may help health students develop skills and encourage them to become accountable health professionals who can support current societal healthcare demands.

References

- Al-Hanawi, M. K. et al. (2020) 'Knowledge, attitude and practice toward COVID-19 among the public in the Kingdom of Saudi Arabia: a cross-sectional study', *Frontiers in public health*. Frontiers, 8, p. 217.
- Albaqawi, H. M. et al. (2021) 'Risk assessment and management among frontline nurses in the context of the COVID-19 virus in the northern region of the Kingdom of Saudi Arabia', *Applied Nursing Research*. Elsevier, 58, p. 151410.
- Biswas, N. et al. (2021) 'The nature and extent of COVID-19 vaccination hesitancy in healthcare workers', *Journal of community health*. Springer, 46(6), pp. 1244–1251.
- Cornwell, C., Mustard, D. B. and Van Parys, J. (2013) 'Noncognitive skills and the gender disparities in test scores and teacher assessments: Evidence from primary school', *Journal of Human resources*. University of Wisconsin Press, 48(1), pp. 236–264.
- Erfani, A. et al. (2020) 'Knowledge, attitude and practice toward the novel coronavirus (COVID-19) outbreak: a population-based survey in Iran', *Bull world Health organ*, 30(10.2471), pp. 10–2471.
- Fontenot, H. B. et al. (2021) 'Undergraduate nursing students' COVID-19 vaccine intentions: A national survey', *Plos one*. Public Library of Science San Francisco, CA USA, 16(12), p. e0261669.
- Grady, D. and Mazzei, P. (2021) 'Doctor 's Death After Covid Vaccine Is Being Investigated', *Health*.
- Islam, M. et al. (2021) 'Knowledge, attitudes and perceptions towards COVID-19 vaccinations: a cross-sectional community survey in Bangladesh', *BMC public health*. Springer, 21(1), pp. 1–11.
- Jiang, N. et al. (2021) 'Nursing students' attitudes, knowledge and willingness of to receive the coronavirus disease vaccine: A cross-sectional study', *Nurse education in practice*. Elsevier, 55, p. 103148.
- Kassianos, G. et al. (2018) 'Motors of influenza vaccination uptake and vaccination advocacy in healthcare workers: A comparative study in six European countries', *Vaccine*. Elsevier, 36(44), pp. 6546–6552.
- Kumar, B., Pinky, S. D. and Nurudden, A. M. (2021) 'Knowledge, attitudes and practices towards COVID-19 guidelines among students in Bangladesh', *Social Sciences & Humanities Open*. Elsevier, 4(1), p. 100194.
- Malik, A. A. et al. (2020) 'Determinants of COVID-19 vaccine acceptance in the US', *EclinicalMedicine*. Elsevier, 26, p. 100495.
- Pasay-An, E. et al. (2021) 'Knowledge, attitudes, and practices regarding COVID-19 among medical and allied health students in the Kingdom of Saudi Arabia', *Journal of Nursing and Midwifery Sciences*. Medknow Publications, 8(2), p. 114.
- Pasay-an, E., Magwilang, J. O. G. and Pangket, P. P. (2020) 'Knowledge, attitudes, and practices of adolescents regarding sexuality and reproductive issues in the Cordillera administrative region of the Philippines', *Makara Journal of Health Research*, 24(3), p. 3.
- Sun, Y. et al. (2020) 'Disease prevention knowledge, anxiety, and professional identity during COVID-19 pandemic in nursing students in Zhengzhou, China', *Journal of Korean Academy of Nursing*. Korean Society of Nursing Science, 50(4), pp. 533–540.
- Szmyd, B. et al. (2021) 'Attitude and behaviors towards SARS-CoV-2 vaccination among healthcare workers: A cross-sectional study from Poland', *Vaccines*. Multidisciplinary Digital Publishing Institute, 9(3), p. 218.

How to cite this article: Jr, R. M., Maestrado, R., Buta, J., Albaqawi, H. M., Rondilla, E., Alsaqri, S., Villareal, S. (2022) 'Knowledge and attitudes toward COVID-19 vaccination among student nurses from Saudi Arabia', *Jurnal Ners*, 17(1), pp. 31-35. doi: <http://dx.doi.org/10.20473/jn.v17i1.34233>

Relationship between stress, anxiety, and depression with suicidal ideation in adolescents

Heni Dwi Windarwati ^{1,*} , Retno Lestari ¹ , Satrio Agung Wicaksono ², Mira Wahyu Kusumawati ¹ , Niken Asih Laras Ati ³ , Shofi Khaqul Ilmy ⁴ , Ari Dwi Sulaksono ¹ , Desi Susanti ⁵

¹ Department of Mental Health Nursing, Faculty of Medicine, Universitas Brawijaya, Indonesia

² Faculty of Computer Science Universitas Brawijaya, Indonesia

³ Saintek Medika Nusantara, Jombang, Indonesia

⁴ Faculty of Medicine, Universitas Pendidikan Ganesha, Indonesia

⁵ School of Nursing, Faculty of Medicine, Universitas Brawijaya, Indonesia

*Correspondence: Heni Dwi Windarwati. Address: Veteran Malang Street 65145, East Java, Indonesia. Email: henipsik.fk@ub.ac.id

Responsible Editor: Ilya Krisnana and Ferry Efendi

Received: 11 August 2021 ◦ Revised: 20 April 2022 ◦ Accepted: 20 April 2022

ABSTRACT

Introduction: Suicide is a significant mental health problem in adolescents worldwide that requires comprehensive treatment. This study examined the relationship between stress, anxiety, and depression with suicidal ideation among Indonesian adolescents.

Methods: This cross-sectional study examined 869 high school adolescents. Suicide ideation was a dependent variable, while stress, anxiety, and depression were independent variables in this study. The instrument used to measure stress, anxiety, and depression is the Depression Anxiety Stress Scale (DASS-21), while the idea of suicide was measured using the Scale for Suicide Ideation (SSI) instrument. Data were analyzed with Spearman statistical test to examine the association between variables.

Results: The results showed that teenagers who had suicidal ideation were 23.7% of respondents, while 12.4% had a high intention of suicidal ideation. A total of 44.4%, 72.7%, and 71.4% of teens experienced stress, anxiety, and depression in the mild to very severe categories, respectively. There was a significant relationship between suicidal ideation with stress levels (p-value < 0.001; r value = 0.267), anxiety (p-value < 0.001; r value = 0.237), and level of depression (p-value < 0.001; r value = 0.233) in adolescents.

Conclusions: The study's findings revealed that stress, anxiety, and depression significantly affect suicide ideation in adolescents. Therefore, to prevent increased suicide ideation in adolescents, mental and psychological health support measures must be given to deal with stress, anxiety, and depression.

Keywords: adolescents; anxiety; depression; stress; suicidal ideation

Introduction

Mental health problems in adolescents have become a worldwide concern. In the last few decades, research on mental health problems in adolescents has been carried out (Bruha et al., 2018), such as in the United States (Mojtabai and Olfson, 2020), UK (Fink et al., 2015), Italy (Frigerio et al., 2009), Australia (Allen and McKenzie, 2015), Korea (Roh et al., 2016), China (Wang, Zhang and Zhang, 2020), and Malaysia (Ahmad et al., 2015). Mental health problems in adolescents are very complex, consisting of bullying, gaming or internet addiction,

emotional disorders (anxiety, depression, frustration, anger), and suicide-self harm (WHO, 2019a).

In 2015, suicide became the 20th most common cause of death globally. Data from WHO (2019) report that around 800,000 people die from committing suicide each year, accounting for 1.5% of all deaths and the second leading cause of death at 15-29 years (WHO, 2019b). In North America, suicide is the 10th most common cause of death (Fazel and Runeson, 2020). In recent years in Indonesia, the phenomenon of suicide has been increasingly worrisome. The WHO estimates that the suicide rate in Indonesia could reach 3.4% of 100,000

people if it does not receive serious attention from various parties (WHO, 2018). Meanwhile, data in East Java province in 2018 showed that as many as 549 suicide cases occurred in East Java (Badan Pusat Statistik, 2019).

Suicide is an action to end life intentionally and is carried out consciously (Wuryaningsih et al., 2018). Suicide consists of various terms, such as self-injury, suicide ideas, suicide plans, and completed suicide. Suicidal ideation is the thought of death caused by oneself. Suicidal ideation can be passive when there are only thoughts of suicide without the intention to act or can be expressed directly, verbally, or not shown at all (Stuart, 2016). Suicidal ideation was found prevalent in adolescents (Strandheim et al., 2014). Someone who has the idea of suicide experiences social isolation, low self-esteem, and lack of trust with others, tends to think negatively, is inflexible, and feels powerless over his life (Austin andKunyk, 2019). Suicidal ideation is positively and significantly related to the occurrence of suicide attempts (Ziaaddini, Navadeh and Saeedi, 2009). So, it is crucial to look at factors related to suicidal ideation.

Among adolescents, the prevalence of suicide attempts, suicide ideation, and self-harm reaches 6%, 18% 13.5%, respectively, where suicide behavior is influenced by internal and external factors such as low individual coping, misuse of smartphones, nutrition imbalance, bad lifestyle, low interaction in family, bullying, and social problems (Lim et al., 2019; Rohmana, Estelina and Iskandar, 2020; Ati, Paraswati and Windarwati, 2021). However, the risk of suicide is also related to attitude toward problems and personality aspects (Sharif et al., 2014), namely hostility, impulsivity, depression, and despair, while anti-social symptoms and depression often appear at a young age (Stuart, 2016). Research related to suicidal ideation among university students in Tehran showed that depression, anxiety, and daily stress were positively associated with suicidal ideation (Izadinia et al., 2010).

Stress is an intrinsic and extrinsic stimulus that causes a biological response and compensatory response to the stimulus (Stewart et al., 2019). Stress can cause various actions on the body, ranging from homeostasis changes to life-threatening effects (Stewart et al., 2019). Exposure to focus is at the core of the theory of suicide, and stress is associated with suicidal ideation, especially in adults and adolescents (Yaribeygi et al., 2017). Anxiety is a normal human emotion, which involves behavior, affective, and cognitive responses when facing the perception of danger. Anxiety is considered excessive or pathological when it causes significant distress or distraction. Anxiety often co-exists with depression (Craske and Stein, 2016). Depression is a common mental disorder characterized by constant sadness and a lack of interest or pleasure in previously beneficial or enjoyable activities (WHO, 2019). Depression is associated with substantial current and future morbidity and increases the risk of suicide (Thapar et al., 2012).

Based on the importance of the risk factors for suicidal ideation in adolescents and the limited research conducted on this topic, especially in Indonesia, the present study aimed to examine the relationship between stress, anxiety, depression, and associated factors with the emergence of suicidal ideation in adolescents. The study's results are expected to be a catalyst in providing healthcare for adolescents with suicidal ideation in Indonesia.

Materials and Methods

Study Design

A cross-sectional survey design was employed in this study.

Respondent

The population and sample were all adolescents aged between 15-16 years from five high schools in East Java Province, Indonesia. The total sampling method was employed to select respondents from five high schools in this study, with sample size of 869 adolescents. The inclusion criteria in this study were 1) high school first-grade adolescents aged 15-16 years, 2) willing to take part in the study, and 3) completing the questionnaire.

Instrument

Data were gathered using a questionnaire to determine the respondents' demographic data, suicide ideation, stress, anxiety, and depression.

Stress, anxiety, and depression were measured using the Depression Anxiety Stress Scale (DASS-21), validated, and translated into the respondents' national language (Indonesian language). DASS-21 is a well-established instrument for measuring depression, anxiety, and stress symptoms in clinical and non-clinical samples (Lovibond and Lovibond, 1995). The instrument consists of 21 questions with an answer rating scale of 0 (never), 1 (sometimes), 2 (often), and 3 (always). The score on DASS-21 will be multiplied by two to calculate the final score. The cut-off score for depression was 0-9 (normal), 10-13 (mild), 14-20 (moderate), 21-27 (high), and > 28 (very high). The cut-off points for anxiety were 0-7 (normal), 8-9 (mild), 10-14 (moderate), 15-19 (high), and > 20 (very high). The cut-off points for stress were 0-14 (normal), 15-18 (mild), 19-25 (moderate), 26-33 (High), and > 34 (Very High) (Lovibond and Lovibond, 1995; Brumby et al., 2011). The DASS-21 Indonesian version questionnaire has passed the validity and reliability test (Cronbach's alpha > 0.7).

Furthermore, suicide ideation was measured using the Scale for Suicide Ideation (SSI) instrument, validated, and translated into the respondents' national language (Indonesian language). The SSI instrument is an ideal tool in the psychological and clinical investigation of suicidal ideation and differentiates individuals based on their level of suicidal ideation (Beck, Kovacs and

Weissman, 1979). The cut-off points for suicide ideation were 0 (did not have suicide ideation), 1 (low/minimal suicide ideation), and >1 (high suicide ideation). The SSI Indonesian version questionnaire has passed the validity and reliability test (Cronbach's alpha > 0.7).

Data Analysis

The data analyses conducted were univariate and bivariate. Descriptive analysis as frequency and percentage were used for categorical variables. Spearman Rank Analysis was used to see the correlation between the two variables. Suicide ideation was a dependent variable, while stress, anxiety, and depression were independent variables in this study. Correlation analysis was said to be significantly related if the p -value < 0.05. Data analysis was performed using Statistical Package for Social Sciences (SPSS) version 16.

Ethical Consideration

This study was approved by the Health Research Ethics Committee of the Health Polytechnic of the Ministry of Health Malang with an Ethical Approval Certificate ("Ethical Approval") Reg. Number. 335 / KEPK-POLKESMA / 2019. Before participating in the research, participants were given an explanation related to the research to be conducted. For ethical reasons, informed consent was sent to the participants. Participants who agreed to participate in this research signed the informed consent. The participation of respondents in this study was voluntary. This study followed ethical research principles, which include beneficence, justice, and respecting human dignity.

Results

Mostly, adolescents in this study did not have suicidal ideation (76.3%). However, as many as 23.7% of adolescents had the intention of suicide ideation, while 12.4% of respondents had high suicide ideation. Most participants did not experience stress (55.6%) or depression (28.5%). In the anxiety variable, it can be seen that the majority of adolescents' experience anxiety in the very high category (36.1%), and as many as 27.3% of adolescents did not have anxiety. In conclusion, 44.4%, 72.7%, and 71.5% of adolescents experienced minimal to very high stress, anxiety, and depression, respectively (Table 1).

Table 2 shows as many as 413 (47.5%) of adolescents who did not intend suicide ideation also did not experience stress. In contrast, 3% of students had a high intention of suicide ideation and very high-stress levels. The bivariate analysis results in Table 2 concluded a significant relationship between stress levels with suicidal ideation in adolescents, as indicated by the p -value of < 0.001. In addition, the correlation coefficient of 0.267 indicated a positive direction between variables.

Table 1 Characteristics of research respondents (N = 869)

Characteristics	n	%
Sex		
Female	426	49.0
Male	443	51.0
Suicidal ideation		
High	108	12.4
Low	98	11.3
None	663	76.3
Stress		
Very high	65	7.5
High	96	11.0
Moderate	131	15.1
Mild	94	10.8
Normal/none	483	55.6
Anxiety		
Very high	314	36.1
High	104	12.0
Moderate	119	13.7
Mild	95	10.9
Normal/none	237	27.3
Depression		
Very high	190	21.9
High	112	12.9
Moderate	205	23.6
Mild	114	13.1
Normal/none	248	28.5

Based on Table 2, the analysis concluded a significant relationship between the level of anxiety with suicidal ideation in adolescents, as indicated by the value of p -value < 0.001. Furthermore, the correlation coefficient of 0.237 indicates a positive relationship between variables.

In terms of depression, it showed a significant relationship between the level of depression with suicidal ideation in adolescents, as indicated by the value of p -value < 0.001. The correlation coefficient of 0.233 shows a positive relationship between variables. This study concluded that stress, anxiety, and depression have a significant and positive relationship with suicide ideation in adolescents. This showed that the lower the stress, anxiety, and depression level, the suicidal ideation in adolescents was also getting lower, and vice versa.

Discussions

The present study investigates the relationship between stress, anxiety, and depression with the emergence of suicidal ideation in adolescents. Based on the findings, this study indicated that stress, anxiety, and depression had a significant and positive relationship with suicide ideation in adolescents. The results of this study were in line with research conducted by Izadinia et al. (2010), which showed that stress, anxiety, and depression correlate positively and significantly with thoughts of suicide. In addition, other studies also indicated that stress, depression, and anxiety were the major risk factors for suicidal behavior (Nguyen et al., 2013; Lew et al., 2019).

The idea of adolescent suicide arises because, at this stage, the emotional development that occurs is still unstable (Asante et al., 2017). The percentage of suicide ideas, suicide plans, and attempted suicides have a high

Table 2 Spearman rank correlation analysis of stress, anxiety, and depression with suicidal ideation

		Suicidal ideation								P-value	r
		High		Low		None		Total			
		n	%	n	%	n	%	n	%		
Stress	Very High	26	3.0	4	0.5	35	4.0	65	7.5	<0.001	0.267
	High	16	1.8	14	1.6	66	7.6	96	11		
	Moderate	25	2.9	26	3.0	80	9.2	131	15.1		
	Mild	15	1.7	10	1.2	69	7.9	94	10.8		
	Normal	26	3.0	44	5.1	413	47.5	483	55.6		
Total		108	12.4	98	11.3	663	76.3	869	100		
		Suicidal ideation								P-value	r
		High		Low		None		Total			
		n	%	n	%	n	%	n	%		
Anxiety	Very High	69	7.9	42	4.8	203	23.4	314	36.1	<0.001	0.237
	High	17	2.0	12	1.4	75	8.6	104	12.0		
	Moderate	5	0.6	12	1.4	102	11.7	119	13.7		
	Mild	7	0.8	14	1.6	74	8.5	95	10.9		
	Normal	10	1.2	18	2.1	209	24.1	237	27.3		
Total		108	12.4	98	11.3	663	76.3	869	100		
		Suicidal ideation								P-value	r
		High		Low		None		Total			
		n	%	n	%	n	%	n	%		
Depression	Very High	49	5.6	25	2.9	116	13.3	190	21.9	<0.001	0.233
	High	14	1.6	18	2.1	80	9.2	112	12.9		
	Moderate	27	3.1	21	2.4	157	18.1	205	23.6		
	Mild	6	0.7	14	1.6	94	10.8	114	13.1		
	Normal	12	1.4	20	2.3	216	24.9	248	28.5		
Total		108	12.4	98	11.3	663	76.3	869	100		

rate at 17 years and 18 years and above (Asante et al., 2017). The percentage of suicide risk rates are high at ages 15 until 18 years, and this age range is a stage of late adolescence (Yeomans and Christensen, 2017), where, in this study, the age of the respondents was between 15-16 years. The previous research stated that stress occurs a lot in adolescence, accompanied by depression and anxiety (Duarte et al., 2019). Youth is a period that often elicits stressful conditions because, during this time, puberty hits as the peak of growth and development, both physically and mentally (Miller and Prinstein, 2019). Stress can occur due to pressure on unwanted thoughts such as the burden or problems inside the family environment (Windarwati et al., 2020). Stress also happens to those who suppress their feelings and refrain from openly expressing emotions (Kim, Bassett and Voisin, 2019). Emotional status in adolescence is still changing between behavior that shows maturity with childish acts. Academic stress factors can play an essential role in determining adolescent mental health (Waghachavare et al., 2013). About 26% of high school students in the study had high levels of educational stress. Stress is a significant risk factor causing depression (Asante et al., 2017). Stress is related to anxiety and depression and is a significant risk factor for teen suicidal ideation.

This study suggested that the lower the stress level, the lower the suicide idea, and vice versa. Various work stressors and related life stressors, such as stressful events, loss, unemployment, and other environmental stressors, link to suicidal ideation (Ibrahim, Amit and Suen, 2014). This study showed that stress was positively related to suicidal ideation. The problem is how to adapt

to stress without experiencing distress because distress can trigger psychological effects such as anxiety and depression. Interaction of various aspects of stress can complicate optimal stress management and generate suicidal ideation (Ibrahim, Amit and Suen, 2014).

The lower the anxiety indicated, the lower the suicidal ideation in adolescents. Adolescents who experience anxiety tend to have a higher level of vigilance and concern about something that has not yet happened. Adolescents who experience anxiety can affect daily activities related to weakness and powerlessness (Craske et al., 2017). Adolescence is a period where individuals are worried about their future and face various pressures from family, school environment, and peers. This situation makes adolescents feel anxious, with emotions that are still unstable, and difficulty in solving difficult problems can make teens have ideas for suicide or self-injury as a way out of anxiety experienced. People with anxiety were more likely to have suicidal ideation, attempted suicide, and completed suicide than those who did not experience anxiety (Malik et al., 2014).

There was a significant relationship between the level of depression with suicidal ideation in adolescents. The lower the depression level indicated, the lower the suicide idea. Depression in adolescents occurs due to overflowing emotions and immaturity in making decisions and continuing the physiological development process. The high incidence of depression is caused by the increased responsibility they have to bear, both responsibility to parents, teachers, and those closest to them. There are various physical, social, emotional, and academic challenges in adolescents. This stage is also the peak age of onset of many mental health disorders (Ibrahim, Amit

and Suen, 2014). Furthermore, psychological conditions of students generally cause depression in adolescents, relationship problems with friends (boy/girlfriend), academic problems at school, relationship problems with parents and family, relationship problems with classmates, financial issues, relationship problems with teachers, and physical health problems (Magklara et al., 2015). Lack of parental supervision, alcohol abuse, and smoking are also significant risk factors for teenage depression (Asante et al., 2017). Suicidal ideation in late teens is caused by depression (Asante et al., 2017). Depression is a significant cause of suicide in adolescents (Towsend, 2018). Depression is associated with feelings of individual despair, helplessness, and lack of social support and skills to overcome difficulties and a stressful life (Ibrahim, Amit and Suen, 2014). Depression can also be a response to the loss of parents, relatives, friends, or the end of a relationship with a lover (Towsend, 2018).

Early adolescence is the initial stage of puberty, the regulation system itself is still immature, so it tends to exhibit unstable emotions (Curtis, 2015). The frontal lobe, which is part of the brain to regulate reasoning and decision-making, begins to develop during early adolescence. Consequently, adolescents tend to act impulsively and think uncritically (UNICEF, 2011). In the late adolescence stage, emotions that are not yet sufficiently stable and a self-regulation system that is still developing gives this age stage an emotional status that is ever-changing between behavior that shows maturity and acting like children (Curtis, 2015). More complex problems in late adolescence lead to a higher prevalence of depression and a higher risk of suicide at this age. However, physical development and growth will continue at this stage, including the frontal lobe, which is part of the brain that governs reasoning and decision-making. Frontal lobes that experience further development cause late adolescents to tend to have the power of reasoning and the ability to make their own decisions well (Curtis, 2015). Therefore, coping mechanisms must be strengthened at this age. Anxiety and depression are related to the idea of suicide and self-injury. Adolescents with suicidal ideation and self-injury tendencies have higher depression, anxiety, and stress levels.

The use of a cross-sectional survey design in the present study is not without limitations. For instance, it cannot determine the causal relationship among variables. However, this study was conducted in a population with a large enough number and the same respondents (10th grade) characteristics, so the results are expected to reduce possible bias. Furthermore, this study can also add to the literature on risk factors for suicide in adolescents in Indonesia, which is still very limited.

Conclusions

This present study indicated that stress levels, anxiety, and depression correlated positively and significantly with suicidal ideation in adolescents. Stress, anxiety, and depression comprise a condition that is closely related to adolescent activities and is close to teenage life. Therefore, nurses have to provide mental health education in the form of counselling, psychoeducation, and screening to adolescents, teachers, and parents of students who are healthy and at-risk of mental health problems. At the same time, students who experience mental health problems that can cause suicide ideas should be given psychotherapy in the form of cognitive-behavioral therapy to overcome their problems. Mental Health Service Providers have to pay great attention to adolescents regarding psychological issues due to emotional burdens experienced by adolescents either at school, at home, or in the environment that can cause mental health problems, especially stress, anxiety, and depression in adolescents. Schools need to develop a school mental health unit to promote and prevent stress in adolescents through early detection, mental health therapy, and psychosocial support. Lastly, further research is needed to identify the causal relationship between stress, anxiety, depression with suicide ideation in Indonesia. It is hoped that a decrease in adolescent suicide ideation has implications for decreasing the incidence of suicide in adolescents.

References

- Ahmad, N. A. et al. (2015) 'Trends and factors associated with mental health problems among children and adolescents in Malaysia', *International Journal of Culture and Mental Health*, 8(2), pp. 125–136. doi: 10.1080/17542863.2014.907326.
- Allen, K. A. and McKenzie, V. L. (2015) 'Adolescent mental health in an Australian context and future interventions', *International Journal of Mental Health*, 44(1), pp. 80–93. doi: 10.1080/00207411.2015.1009780.
- Asante, K. O. et al. (2017) 'The prevalence and correlates of suicidal behaviours (ideation, plan and attempt) among adolescents in senior high schools in Ghana', *SSM - Population Health*, 3(May), pp. 427–434. doi: 10.1016/j.ssmph.2017.05.005.
- Ati, N. A. L., Paraswati, M. D. and Windarwati, H. D. (2021) 'What are the risk factors and protective factors of suicidal behavior in adolescents? A systematic review', *Journal of Child and Adolescent Psychiatric Nursing*, 34(7), pp. 7–18. doi: 10.1111/jcap.12295.
- Austin, W & Kunyk, D. (2019) *Psychiatric and Mental Health Nursing for Canadian Practice*. Cina: Wolters Kluwer.
- Badan Pusat Statistik (2019) *Banyaknya Desa/Kelurahan Menurut Keberadaan Korban Bunuh Diri, Lokasi Berkumpul Anak Jalanan, Gelandangan, dan Pekerja Seks Komersial (PSK)*, 2018. Jawa Timur.
- Beck, A. T., Kovacs, M. and Weissman, A. (1979) 'Assessment of suicidal intention: The Scale for Suicide Ideation', *Journal of Consulting and Clinical Psychology*, 47(2), pp. 343–352. doi: 10.1037//0022-006x.47.2.343.
- Bruha, L. et al. (2018) 'Global child and adolescent mental health: challenges and advances', *London Journal of Primary Care*, 10(4), pp. 108–109. doi: 10.1080/17571472.2018.1484332.
- Brumby, S. et al. (2011) 'Reducing psychological distress and obesity in Australian farmers by promoting physical activity', *BMC Public Health*, 11(1), p. 362. doi: 10.1186/1471-2458-11-362.
- Craske, M. G. et al. (2017) 'Anxiety disorders', *Nature Reviews Disease Primers*, 3. doi: 10.1038/nrdp.2017.24.
- Craske, M. G. and Stein, M. B. (2016) 'Anxiety', *The Lancet*, 388(10063), pp. 3048–3059. doi: 10.1016/S0140-6736(16)30381-6.
- Curtis, A. C. (2015) 'Defining the adolescent', *Journal of Adolescent and Family Health*, 7(2). Available at: <https://scholar.utc.edu/jafh/vol7/iss2/2>.

- Duarte, T. A. et al. (2019) 'Self-harm as a predisposition for suicide attempts: A study of adolescents' deliberate self-harm, suicidal ideation, and suicide attempts', *Psychiatry Research*, 287. doi: 10.1016/j.psychres.2019.112553.
- Fazel, S. and Runeson, B. (2020) 'Suicide', *New England Journal of Medicine*, 382(3), pp. 266–274. doi: 10.1056/NEJMra1902944.
- Fink, E. et al. (2015) 'Mental health difficulties in early adolescence: A comparison of two cross-sectional studies in England from 2009 to 2014', *Journal of Adolescent Health*, 56(5), pp. 502–507. doi: 10.1016/j.jadohealth.2015.01.023.
- Frigerio, A. et al. (2009) 'Prevalence and correlates of mental disorders among adolescents in Italy: The PRISMA study', *European Child and Adolescent Psychiatry*, 18(4), pp. 217–226. doi: 10.1007/s00787-008-0720-x.
- Ibrahim, N., Amit, N. and Suen, M. W. Y. (2014) 'Psychological factors as predictors of suicidal ideation among adolescents in Malaysia', *PLoS ONE*, 9(10). doi: 10.1371/journal.pone.0110670.
- Izadinia, N. et al. (2010) 'A study of relationship between suicidal ideas, depression, anxiety, resiliency, daily stresses and mental health among Tehran university students', *Procedia - Social and Behavioral Sciences*, 5, pp. 1615–1619. doi: 10.1016/j.sbspro.2010.07.335.
- Kim, D. H., Bassett, S. M. and Voisin, D. R. (2019) 'Family stress and youth mental health problems: Self-efficacy and future orientation mediation', *American Journal of Orthopsychiatry*, 89(2), 125. doi: <https://doi.org/10.1037/ort0000371>.
- Lew, B. et al. (2019) 'Associations between depression, anxiety, stress, hopelessness, subjective well-being, coping styles and suicide in Chinese university students', *PLoS ONE*, 14(7), pp. 1–10. doi: 10.1371/journal.pone.0217372.
- Lim, K. S. et al. (2019) 'Global lifetime and 12-month prevalence of suicidal behavior, deliberate self-harm and non-suicidal self-injury in children and adolescents between 1989 and 2018: A meta-analysis', *International Journal of Environmental Research and Public Health*, 16(22). doi: 10.3390/ijerph16224581.
- Lovibond, P. and Lovibond, S. (1995) *Manual for the Depression Anxiety Stress Scales*. 2nd edn. Sydney: Psychology Foundation.
- Magklara, K. et al. (2015) 'Depression in late adolescence: A cross-sectional study in senior high schools in Greece', *BMC Psychiatry*, 15(1). doi: 10.1186/s12888-015-0584-9.
- Malik, S. et al. (2014) 'The association between sleep disturbances and suicidal behaviors in patients with psychiatric diagnoses: A systematic review and meta-analysis', *Systematic Reviews*. doi: 10.1186/2046-4053-3-18.
- Miller, A. B. and Prinstein, M. J. (2019) 'Adolescent Suicide as a Failure of Acute Stress-Response Systems', *Annual Review of Clinical Psychology*, 15(1), pp. 425–450. doi: 10.1146/annurev-clinpsy-050718-095625.
- Mojtabai, R. and Olfson, M. (2020) 'National Trends in Mental Health Care for US Adolescents', *JAMA Psychiatry*, 77(7), pp. 703–714. doi: 10.1001/jamapsychiatry.2020.0279.
- Nguyen, D. T. et al. (2013) 'Depression, anxiety, and suicidal ideation among Vietnamese secondary school students and proposed solutions: A cross-sectional study', *BMC Public Health*, 13(1). doi: 10.1186/1471-2458-13-1195.
- Roh, S. et al. (2016) 'Mental health services and R&D in South Korea', *International Journal of Mental Health Systems*, 10(1), pp. 1–10. doi: 10.1186/s13033-016-0077-3.
- Rohmana, D. Y., Estelina, K. and Iskandar, I. (2020) 'The Bullying Phenomenon and Handling Efforts in Reducing Cases of Bullying: A Systematic Review', *Jurnal Ners*, 15(2). doi: <http://dx.doi.org/10.20473/jn.v15i2.20479> This.
- Sharif, F. et al. (2014) 'Comparison of Personality Traits, Coping Styles, and Psychiatric Disorders in Adult Suicidal and Non-Suicidal Individuals', *International journal of community based nursing and midwifery*, 2(3), pp. 148–156.
- Stewart, J. G. et al. (2019) 'Life Stress and Suicide in Adolescents', *Journal of Abnormal Child Psychology*, 47(10), pp. 1707–1722. doi: 10.1007/s10802-019-00534-5.
- Strandheim, A. et al. (2014) 'Risk factors for suicidal thoughts in adolescence—a prospective cohort study: The Young-HUNT study', *BMJ Open*, 4(8). doi: 10.1136/bmjopen-2014-005867.
- Stuart, G. . (2016) *Prinsip dan praktik keperawatan jiwa Stuart (terjemahan) Jilid 1 & 2*. Singapore: Elsevier.
- Thapar, A. et al. (2012) 'Depression in adolescence', *Lancet*, 379(9820), pp. 1056–1067. doi: 10.1016/S0140-6736(11)60871-4.
- Towsend, M. C. (2018) *Psychiatric mental health nursing: Concepts of care in evidence-based practice*. Philadelphia: F. A. Davis Company.
- UNICEF (2011) *Adolescence An Age of Opportunity*.
- Waghachavare, V. B. et al. (2013) 'A Study of Stress among Students of Professional Colleges from an Urban area in India', *Sultan Qaboos University Medical Journal*, 13(August), pp. 429–436.
- Wang, C., Zhang, P. and Zhang, N. (2020) 'Adolescent mental health in China requires more attention', *The Lancet Public Health*, 5(12), p. e637. doi: 10.1016/S2468-2667(20)30094-3.
- WHO (2018) *Suicide rate estimates, crude Estimates by country*, Global Health Observatory data repository. Available at: <https://apps.who.int/gho/data/node.main.MHSUICIDE> (Accessed: 23 April 2020).
- WHO (2019a) *Adolescent mental health, Fact sheets*. Available at: <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health> (Accessed: 28 August 2020).
- WHO (2019b) *Fact Sheets: Suicide*. Available at: <http://www.who.int/en/news-room/fact-sheets/detail/suicide> (Accessed: 20 August 2020).
- WHO (World Health Organization) (2019) *Depression, Health topics*. Available at: https://www.who.int/health-topics/depression#tab=tab_1 (Accessed: 17 October 2020).
- Windarwati, H. D. et al. (2020) 'The Relationship between Family Harmony with Stress, Anxiety, and Depression in Adolescents', *Jurnal Ners*, 15(3). doi: <http://dx.doi.org/10.20473/jn.v15i2.21495>.
- Wuryaningsih, E. W. et al. (2018) *Keperawatan Kesehatan Jiwa*. Jember: Percetakan & penerbitan Universitas Jember.
- Yaribeygi, H. et al. (2017) 'The impact of stress on body function: A review', *EXCLI Journal*, 16, pp. 1057–1072. doi: 10.17179/excli2017-480.
- Yeomans, C. and Christensen, H. (2017) *Youth Mental Health Report Youth Survey 2012-16*.
- Ziaaddini, H., Navadeh, S. and Saeedi, A. (2009) 'Prevalence of Suicide Ideation, Attempts and the Associated Factors among a Sample of Iranian Population in South Part of the Country: A Population Based Study', *Iranian Journal of Psychiatry*, 4(3), pp. 92–96.

How to cite this article: Windarwati, H. N., Lestari, R., Wicaksono, S. A., Kusumawati, M. W., Ati, N. A. L., Ilmy, S. K., Sulaksono, A. D., Susanti, D. (2022) 'Relationship between stress, anxiety, and depression with suicidal ideation in adolescents', *Jurnal Ners*, 17(1), pp. 36–41. doi: <http://dx.doi.org/10.20473/jn.v17i1.31216>

The correlation between parental communication pattern, self-esteem, and moral disengagement with cyberbullying behavior in early adolescents: A cross-sectional study

Dian Octavia ^{1,*} , Rian Maylina Sari ¹ , Diah Merdekawati ¹ , Rara Marisdayana ¹ , Rian Yuliyana ² 

¹ Institute of Health Science Harapan Ibu Jambi, Indonesia

² Health Polytechnic Tanjung Pinang, Indonesia

*Correspondence: Dian Octavia. Address: Kol. Tarmizi Kodir Street No.71, Pakuan Baru, Jambi City, 36122, Indonesia.
 Email: octaviadian04@gmail.com

Responsible Editor: Ilya Krisnana

Received: 12 January 2021 ◦ Revised: 9 October 2021 ◦ Accepted: 13 October 2021

ABSTRACT

Introduction: Cyberbullying has become a new case in young millennials that can lead to being a serious problem if it is not faced properly. Many cases cannot be resolved because the victims did not tell the truth clearly to their parents and teachers. It may cause a lack of self-esteem among the students and also make the perpetrator become morally disengaged. The purpose of this study was to identify the correlation between parental communication patterns, moral disengagement, and self-esteem with cyberbullying behavior in junior high school.

Methods: A cross-sectional study was performed with 362 junior high school students in Indonesia using purposive sampling. Variables in this study were moral disengagement, self-esteem and parental communication patterns with cyberbullying behavior. The data were collected with the Rosenberg self-esteem scale, the cyberbullying behavior scale, the moral disengagement scale and the family communication pattern scale. Bivariate analysis with Chi-square was used to analyze the data.

Results: This study found out that there were 223 students, 61.6% were female, and 43.6% were aged 14 years. There were 187 students (51.7%) who had mild cyberbullying behavior. The study revealed that there were significant correlations among cyberbullying behavior with moral disengagement ($p = 0.005$), self-esteem ($p = 0.008$), and parental communication pattern ($p = 0.019$).

Conclusions: Parental communication, moral disengagement and self-esteem have a correlation with cyberbullying behavior. It was expected that the schools need to implement a training to improve social skills in junior high school students.

Keywords: cyberbullying; moral disengagement; parental communication pattern; self-esteem

Introduction

The development of communication and information technology also has great impact on the development of today's generation, which we know as millennials society (society 5.0 era). Convenience in almost all daily activities can now be accessed very easily everywhere and any time. But, besides those great impacts, we cannot be blind to one of the negative impacts which be experienced by our young generation, that is cyberbullying.

Cyberbullying is a type of bullying which uses digital technologies and take place on social media, messaging

platforms, gaming platforms or websites (Febriana and Fajrianti, [2019](#)). Practically, it has many negative impacts on victims such as depression low self-esteem, difficulty concentrating in class, anxiety and suicide (Talei and Ardani, [2009](#); Patchin and Hinduja, [2012](#); Rahayu, [2012](#)). Other studies have found that bullying causes stress, emotional, social anxiety, and risk of drug use (Sartana and Afriyeni, [2017](#)). Another study revealed that students who experienced bullying in cyberspace were twice as likely to experience adverse health effects, such as headaches and stomachache, than those who did

not experience bullying (Nursalam et al., 2019). In addition, another study stated that only one out of three adolescents who had been victims of cyberbullying shared their experiences with their parents, teachers and friends (Mendez-Baldwin et al., 2015). It shown that many cases are still uncovered and victims did not receive proper intervention to recover from the situation.

The Indonesian Child Protection Commission (KPAI) stated child victims of cybercrime in Indonesia reached 679 cases and are growing very rapidly (KPAI, 2019). The prevalence among junior high school students was around 48.2% (Safaria, 2016). One factor that makes a child becomes of cyberbullying perpetrator is a grudge against the victim (Pandie and Weismann, 2016). There are several factors influencing the motives of cyberbullying behavior, namely family factors, failure to control oneself, and environmental factors. The poor quality of communication between parents and adolescents can cause adolescent behavior deviations (Nursalam et al., 2019). Other research also stated that through positive communication and interaction within the family, parents and adolescents can strengthen interpersonal relationships, so that adolescents would not be easily influenced by negative invitations, including becoming bullies, by their friends (Firdanianty, Lubis, et al., 2016).

Based on the preliminary survey in four junior high schools in four different districts of Jambi City, Indonesia, it was found that junior high school students tend not to open up with their parents if they become victims of bullying in social media. Students are also reluctant to report the incident to their teachers and tell their close friends about problems instead. When it comes to asking about committing cyberbullying, they seemed to be reluctant to tell the truth. They said that they tend to be victims of bullying in social media. Considering the lack of research related to cyberbullying in Indonesia, this study aimed to identify the correlation among parental communication patterns, moral disengagement, and self-esteem with cyberbullying behavior in junior high schools in Jambi City.

Materials and Methods

Study Design

This was a quantitative study with a cross sectional design which aimed at determining the tendency of the early adolescents with cyberbullying behavior in junior high schools. The study sample was selected by purposive sampling technique. The study location was chosen by multistage random sampling. The study was carried out in two months between October and November, 2020. The independent variables were parental communication patterns, moral disengagement, and self-esteem, while the dependent variable was cyberbullying behavior.

Respondents

The population of this study was all students from four selected junior high schools in Jambi City, with 6,541 students in total. The sample size was 362 students selected by purposive sampling methods and calculated using Lemeshow formula (Sopiyudi Dahlan, 2016). The samples were divided into four junior high schools using proportional sampling technique. The inclusion criteria were students registered as first, second and third year in selected public junior high schools in four districts (Kota Baru District, represented by Public Junior High School 14; Telanaipura District, Public Junior High School 17; East Jambi District, Public Junior High School 9, and Paal Merah District, Public Junior High School 4). The students must be able to communicate and willing to participate. The students will be excluded if they did not attend school during the data collection process.

Instrument

Cyberbullying Behavior Questionnaire

The data were collected using Cyberbullying Behavior Questionnaire (CBQ), which consists of 20 items covering a wide range of work-related cyberbullying behaviors (Jonsson, Muhonen and Backstrom, 2017). The participants scored the items on a four-point scale, 1 = never, 2 = monthly, 3 = weekly, 4 = daily. The scale has an internal consistency with a coefficient alpha of 0.638 to 0.879 (Husna, Tentama and Purwadi, 2020).

Table 1 Demographic characteristics of the respondents (N = 362)

Variables	n	%
Genders		
Male	139	38.4
Female	223	61.6
Age (years)		
11	8	2.2
12	77	21.3
13	101	27.9
14	158	43.6
15	17	4.7
16	1	3
Cyberbullying information		
Social media	252	69.6
Friends	60	16.57
Teacher	15	4.1
Parents	10	2.76
Not know	25	6.9
Parents' education levels		
Primary school	29	8.0
Junior high school	86	23.75
Senior high School	197	54.1
Bachelor high school	50	13.8
Family Income		
≥ Rp 1.750.000,-	150	41.4
≤ Rp 1.750.000,-	212	58.5

Table 2 Relationship between parental communication pattern, self-esteem, and moral disengagement with cyberbullying behavior (N = 362)

Variable	Cyberbullying behavior				Total		p-value
	Mild		Strong		n	%	
	n	%	n	%			
Parental Communication							
Bad	86	46	103	58.9	189	52.2	0.019
Good	101	54	72	41.1	173	47.8	
Self-Esteem							
Low	86	46	101	57.7	187	51.7	0.008
High	101	54	74	42.3	175	48.3	
Moral Disengagement							
Mild	110	58.8	76	43.4	186	51.4	0.005
Strong	77	41.2	59	56.6	176	48.6	
TOTAL	187	100	175	100	362	100	

Rosenberg Self-Esteem Scale

Rosenberg Self-Esteem Scale (RSES) was developed by Rosenberg in 1965. It consists of 10 items that refer to self-respect and self-acceptance rated on a four-point Likert-scale, from 1 (totally disagree) to 4 (totally agree). Five items were worded positively (i.e., 1, 3, 4, 7, and 10) and five items negatively (i.e., 2, 5, 6, 8, and 9) (Martin-Albo, Nunez and Navarro, 2007). The scale has an internal consistency with a coefficient alpha of 0.40 to 0.97 (Tinakon and Nahathai, 2012).

Moral Disengagement Scale

Moral disengagement constructs include vilifying victims through attribution of blame, reinterpreting conduct or moral justification, hiding personal causal responsibility, and misrepresenting the consequences of individual action (Bussey, Fitzpatrick and Raman, 2015). The participants in this study were asked how much they agreed, starting from 1 = strongly disagree, 2 = disagree, 3= agree, 4 =strongly agree). The scale of internal consistency found a coefficient alpha of 0.80 to 0.87 (Whang and Yang, 2010).

Family communication pattern

The family communication pattern consisted of 28-item Likert scale (Gupta, 2019). This study used the four-point scale, namely 1 = strongly disagree; 2 = disagree; 3= agree; 4 = strongly agree. The scale of internal consistency had a Cronbach’s alpha of 0.869.

The validity and reliability of the test were carried out at a public junior high school in Jambi City on August 8, 2020, with the total of 67 question items. The validity value with the RSES was 0.87. The CBQ was on a scale of 0.98, the moral disengagement scale was at 0.87, and the parental communication pattern scale was at 0.85. The reliability test of the overall instruments, cyberbullying behavior, self-esteem, moral disengagements and family communication pattern, were 0.88, 0.86, 0.83, and 0.81, respectively.

Data Analysis

The data were analysed using univariate analysis to describe the demographic characteristics, cyberbullying

behavior and all independent variables (i.e., self-esteem, moral disengagement and parental communication pattern). The normality data was tested by Kolmogorov Smirnov test for large sample. It was found that the p-value were under 0.05 for all variables, which means the data were not normal (Sopiyudi Dahlan, 2016). The data used median to categorize all variables. The bivariate correlation used Chi-square test to analyze independent and dependent variables. The correlation coefficients varied from -1 to +1. This study indicates a perfectly positive linear relationship. For a positive coefficient, as one variable increases, the other also increases. The analysis found strong relationship between independent and dependent variables in which $\alpha = 0.05$ was used for analysis. The data analysis was processed in Statistical Package for the Social Sciences (SPSS file version 1.0.0-51). The data analysis included data screening and assumption of primary data analysis.

Table 3 The types of cyberbullying behaviors in early adolescence (N = 362)

Type of cyberbullying	n	%
Flaming		
Low (6-9)	194	53.6
High (10-24)	168	46.4
Harassment		
Low (2)	215	59.4
High (3-8)	147	40.6
Denigration (put downs)		
Low (3-4)	245	67.7
High (5-12)	117	32.3
Cyber-stalking		
Low (2)	302	83.4
High (3-8)	60	16.6
Impersonation		
Low (1)	312	86.2
High (2-5)	50	13.8
Outing		
Low (2)	198	54.7
High(3-10)	164	45.3
Exclusion		
Low (1)	239	66.0
High (2-5)	123	34.0
Cyber-threat		
Low (3)	291	80.6
High (4-15)	71	19.4
Anonymity		
Low (3)	290	80.1
High (4-15)	72	19.9

Ethical Consideration

This research has got ethical approval with number: LB.02.06/2/123/2020. The ethical review was tested for all aspects in this research, including scope of study, methodology and the questionnaire. The researchers applied ethics principles of anonymity / confidentiality, beneficence and non-maleficence, autonomy, and justice. An explanation of the background and aim of the study, as well as informed consent, were contained together with the instrument.

Results

The total of 362 early adolescents registered in four junior high schools in Jambi City based on several sub-districts were recruited. The characteristics of early adolescents showed that the majority 223 (61.60%) were female, and almost a half, 158, were aged 14 years old (43.60%; see [Table 1](#)).

[Table 2](#) shows that parental communication pattern, self-esteem, and moral disengagement have significant correlation with cyberbullying behavior of students at the four public junior high schools. All p-value were below 0.05; parental communication ($p = 0.019$); self-esteem ($p = 0.008$); and moral disengagement ($p = 0.005$).

[Table 3](#) shows that all cyberbullying types were in low categories, but high categories in few variables nearly reached a half, namely flaming ($n = 168$; 46.4%), outing ($n = 164$; 45.3%), and harassment ($n = 147$; 40.6%).

Discussions

The majority of respondents were female and almost a half of 158 were at the age of 14 years old. Boys and girls both participate in cyberbullying, although for different reasons. They also use different methods. Girls tend to use a more passive approach, such as spreading rumors and reputation and relationship damage. Boys tend to use direct and cyber-threats intended to revenge. In addition, another view says that, in carrying out their actions, women are more often the target of cyberbullying, while men tend to be the main perpetrators of violence in cyberspace (Pandie and Weismann, [2016](#)).

This study found there were associations between self-esteem, parental communication pattern, and moral disengagement with cyberbullying behavior. Several factors that could influence the variables were friends, parents and school environment, parents' educational levels, and parents' salary (Aini, [2018](#)).

Self-esteem was a component of self-concept factors that can influence self-concept such as pressure from outside from good peers (Yuliayana and Wichaikull, [2014](#)). A peer group was a group of friends who had strong emotional ties and students who could be accepted, socialized, exchanged ideas, and experiences in providing the change and the development of social life. The experience of being a victim can lower person's self-

esteem, or people with low self-esteem are more likely to be targeted as victims.

The quality of communication between parents and adolescents should be improved so the adolescents could get guidance and education to avoid cyberbullying. The poor quality of communications between parents and adolescents can cause adolescent behavior deviations (Gunawan, [2013](#)). Other research stated communication between parents and adolescents that is less than optimal will cause adolescents to be easily influenced by deviant behavior (Luk et al. [2010](#)). Positive communication and interaction within family, parents and adolescents can strengthen interpersonal relationships, so adolescents will not be easily influenced by invitations to do negative activities from their friends (Firdianity, Djua et al., [2016](#)).

Reframing behavior is accomplished through moral justifications, euphemistic labelling, and advantageous comparisons, which enable the individual to view their immoral behavior as ultimately moral or benign. Individuals could also morally disengage by changing their perceptions of their own personal responsibility for the behavior by displacing or diffusing responsibilities to others. Moral disengagement mechanisms blame the target for their suffering, or dehumanize the target, act to reframe the individual's perceptions about the target's role for the behavior. Finally, the harmful outcomes of the immoral behavior can be minimized or viewed as providing benefits for the victim (Bandura, [2002](#)). Cyber-communication may promote hostile attributions of intent, and triggering attributions of blame (Runions, [2013](#)). Ambiguous communications, that are common on social media, may provoke self-justifications of cyber-aggression as retaliatory responses, with responsibility for the perceived provocation attributed to the other person; one's own behavior is perceived as merely a reaction. These processes may be particularly pertinent to aggressive-victims (i.e., bully-victims), whose responses to perceived provocation fuel cycles of bullying (Salmivalli and Nieminen, [2001](#)).

This study also found most adolescents have mild moral disengagement, self-esteem and cyberbullying behavior. This study was in line with Jiang et al. ([2021](#)), that stated adolescents whose information was on the topic of self-acceptance and competence, or having good self-esteem, could judge themselves better than most people. According to other research, the sources related to bullying and self-esteem have consistently found that victims of bullying tend to have lower self-esteem than non-victims (Patchin and Hinduja, [2012](#)). Among people with different levels of self-esteem, victims of bullying tend to have lower self-esteem than non-victims.

This study has several limitations. It only identified the correlation between independent and dependent variables with cross-sectional approach that cannot show causality. This study also did not examine any intervention to increase self-esteem, improve family

communication, address moral disengagement, or prevent cyberbullying behavior. Therefore, further study is needed to address these aspects.

Conclusions

There is a correlation between self-esteem, moral disengagement, parental communication pattern and cyberbullying behavior in early adolescence in junior high school. Further research need to address factors related to moral disengagement between genders, improvement by social group about self-esteem, and disengagement in cyberbullying behavior.

References

Aini, D. F. N. (2018) 'Self in elementary school age children for prevention of bullying cases', *Elementary School of Thought and Development*, 6(1), pp. 36–46. doi: <https://doi.org/10.22219/jp2sd.v6i1.5901>.

Bandura, A. (2002) 'Selective Moral Disengagement in The Exercise of Moral Agency', *Journal of Moral Education*, 31, pp.101–119.

Bussey, K., Fitzpatrick, S. and Raman, A. (2015) 'The role of moral disengagement and self-efficacy in cyberbullying.', *Journal of School Violence*, 14(1), pp. 30–46. doi: 1080/15388220.2014.954045.

Febriana, S. K. T. and Fajrianti (2019) 'Cyber Incivility Perpetrator: The Influenced of Dissociative Anonymity, Invisibility, Asynchronicity, and Dissociative Imagination', in *1st International Conference on Advance and Scientific Innovation, ICASI 2018*. Universitas Lambung Mangkurat, Banjarmasin, Indonesia: Institute of Physics Publishing. doi: 10.1088/1742-6596/1175/1/012238.

Firdanianty, Djuara, L. P., et al. (2016) 'Adolescent Communication Patterns and Influence on Emotional Intelligence of High School Students in Bogor City', *Coommunication Journal of Indonesian Association of Communication of Scholars*, 01, pp. 37–47.

Gunawan, H. (2013) 'Jenis pola komunikasi orang tua dengan anak perokok aktif di Desa Jembayan Kecamatan Loa Kulu Kabupaten Kutai Kartanegara', *eJournal Ilmu Komunikasi*, 1(3), pp. 1–5.

Husna, M., Tentama, F. and Purwadi (2020) 'Validity and Reliability Analysis of the Cyberbullying Scale', *American Scientific Research Journal for Engineering, Technology, and Sciences (ASRJETS)*, 71(1), pp. 20–33.

Jiang, S. et al. (2021) 'Cyber victimization and psychological well-being among Chinese adolescents: Mediating role of basic psychological needs satisfaction and moderating role of positive parenting', *Children and Youth Services Review*. Elsevier, 130, p. 106248.

Jonsson, S., Muhonen, T. and Backstrom, R. C. F. M. (2017) 'Assessing Exposure to Bullying through Digital Devices in Working Life: Two Versions of a Cyberbullying Questionnaire (CBQ)', *Psychology*, 8(3), pp. 477–494. doi: 10.4236/psych.2017.83030.

Jose Martin-Albo, Juan L. Nunez, Jose G. Navarro, and F. G. (2007) 'The Rosenberg Self-Esteem Scale: Translation and Validation in University Students', *The Spanish Journal of Psychology*, 10(2), pp. 458–467. doi: 10.1017/S1138741600006727.

KPAI (2019) 'Catatan KPAI di Hardiknas. Kasus Anak Bully Guru Meningkatkan Drastis _ Komisi Perlindungan Anak Indonesia (KPAI)', 24 July. Available at: <https://www.kpai.go.id/publikasi/kpai-sebut-anak-korban-kejahatan-dunia-maya-capai-679-kasus>.

Luk, J.W., Farhat, T., Iannotti, R.J., & Morton, B. G. (2010) 'Parent-child communication and substance use among adolescents: do father

and mother communication play a different role for sons and daughters?', *Addictive Behaviors*, 35:426–431, 35, pp. 426–431.

Mendez-Baldwin, M. et al. (2015) 'Cyberbullying Among Teens', *Journal of Bullying and Social Aggression*, 1(1).

Nursalam, N. et al. (2019) 'Association between insomnia and social network site use in Indonesian adolescents', *Nursing Forum*. Department of Mental Health, Gerontology and Community Nursing, Faculty of Nursing, Universitas Airlangga, Surabaya, Indonesia: John Wiley and Sons Inc., 54(2), pp. 149–156. doi: 10.1111/nuf.12308.

Pandie, M. M. and Weismann, I. T. J. (2016) 'The Effect of Cyberbullying on Social Media on Reactive Behavior as Perpetrators and Victims of Cyberbullying in Christian Middle School Students in Makassar', *Jurnal Jaffray*, 14(1), pp. 42–62.

Patchin, J. W., & Hinduja, S. (2012) 'Cyberbullying Prevention and Respons', in *Cyberbullying Prevention and Respons*. New York: Routledge.

Rahayu, flourensia sapy (2012) 'Cyberbullying as a Negative Impact of Information Technology Users. Journal of Information Systems', *Journal of informations Systems*, 8(1). doi: <https://doi.org/10.21609/jsi.v8i1.321>.

Runions, K. C. (2013) 'Toward a Conceptual Model of Motive and Self-Control in Cyber-Aggression: Rage, Revenge, Reward, and Recreation', *Journal of & Youth Adolescence*, 42, 751–771, 42(5), pp. 751–771. doi: 10.1007/s10964-013-9936-2.

Safaria, T. (2016) 'Prevalence and impact of cyberbullying in a sample of Indonesian junior high school students.', *Turkish Online Journal of Educational Technology-TOJET*. ERIC, 15(1), pp. 82–91.

Salmivalli, C. and Nieminen, E. (2001) 'Proactive and reactive aggression among school bullies, victims, and bully-victims', *Aggressive Behavior*, 28(1), pp. 30–44. doi: <https://doi.org/10.1002/ab.90004>.

Sartana and Afriyeni, N. (2017) 'Cyber Bullying in Early Adolescence. Journal of Psychology Insight', *Jurnal Psikologi Isight*, 1(1), pp. 25–39.

Savita Gupta, G. (2019) 'Family Communication Patterns Questionnaire: Development and Validation', *International Journal of Recent Technology and Engineering (IJRTE)*, 8(IC2), pp. 542–550.

Sopiyudi Dahlan, M. (2016) *Sample size in medical and health research, series 2 edition 4*. 2nd edn. Jakarta: Epidemiolog Indonesia.

Sopiyudi Dahlan, M. (2021) *Statistics for Medicine and Health*. Indonesia: Salemba Medika.

Talaei, A., B, M. R. F. and Ardani, A. R. (2009) 'Gender, Self-Esteem, Social Support and Depression Among Seventh Grade Students in Junior High Schools', *Iranian Journal of Psikiatry*, 4, pp. 17–22.

Tinakon, W. and Nahathai, W. (2012) 'A Comparison of Reliability and Construct Validity between the Original and Revised Versions of the Rosenberg Self-Esteem Scale', *Psychiatric Symptomatology*, 9(1), pp. 54–58. doi: <https://doi.org/10.4306/pi.2012.9.1.54>.

Whang, xhing-chao and Yang, J. (2010) 'Reliability and validity of Moral Disengagement Scale in Chinese students', *Chinese Journal of Clinical Psychology*, 18(2), pp. 177–179.

Yuliyana, R. and Wichaikull, S. (2014) 'Gender, Self-Esteem, Social Support and Depression Among Seventh Grade Students in Junior High Schools', *International Proceedings of Social and Behavioral Sciences*, 2(1), pp. 142–148.

How to cite this article: Octavia D., Sari R. M., Merdekawati D., Marisdayana R., Yuliyana R. (2022) 'The correlation between parental communication pattern, self-esteem, moral disengagement with cyberbullying behavior in early adolescents', *Jurnal Ners*, 17(1), pp. 42-46. doi: <http://dx.doi.org/10.20473/jn.v17i1.24539>

The development of assessment instrument for postpartum patients with severe preeclampsia-eclampsia based on need for help and self-care models

Endah Suprihatin ^{1,*} , Sri Hardi Wuryaningsih ² 

¹ Health Polytechnic of the Ministry of Health, Surabaya, Indonesia

*Correspondence: Endah Suprihatin. Address: Pucang Jajar Tengah Street No.56, Kertajaya, Surabaya City, East Java 60282, Indonesia. Email: endah.atin@poltekkesdepkes-sby.ac.id

Responsible Editor: Retnayu Pradanie

Received: 15 April 2022 ◦ Revised: 25 April 2022 ◦ Accepted: 26 April 2022

ABSTRACT

Introduction: One of the causes of the high mortality rate of patients with severe preeclampsia-eclampsia in the postpartum period is the inaccurate assessment of the patient's need for help and independence in self-care. This study aimed to develop specific assessment instruments for postpartum patients based on Wiedenbach's need for help and Orem's self-care models.

Methods: This study used a research and development (R&D) approach. The samples were 30 midwives and 100 documents of postpartum patients with preeclampsia from Airlangga University Hospital, Surabaya, and 30 midwives from Haji Hospital Surabaya. Data were collected through document tracking and interviews. Data were analyzed with R&D methods.

Results: The results showed that the mean rank value of the new assessment instrument was higher (29.72) than the old assessment instrument (19.93). A significant difference was found between the new assessment instrument and the old assessment instrument (Wilcoxon test value -5.379 with $p < 0.001$). The new instrument had functionality, efficiency, and usability for assessment of the need for help and self-care of postpartum patients with severe preeclampsia and eclampsia.

Conclusions: Postpartum patients with severe preeclampsia-eclampsia should be assessed by the specific assessment instruments based on Wiedenbach's need for help and Orem's self-care models.

Keywords: assessment instruments; eclampsia; need for help; postpartum; self-care; severe preeclampsia

Introduction

Preeclampsia and eclampsia are one of the main causes of maternal and fetal mortality and morbidity in addition to bleeding and infection (triad complications) (Shamsi, Saleem and Nishter, 2013). In 2019, the maternal mortality rate of East Java Province reached 89.81 per 100,000 live births. The most common cause of maternal death in Surabaya is preeclampsia-eclampsia, which is 26.90% (Ministry of Health, 2021).

Preeclampsia is a widespread disorder of endothelial or vascular malfunction resulting in vasospasm after 20 weeks of gestation, resulting in decreased organ perfusion and endothelial activation, leading to hypertension, non-dependent edema, and proteinuria. It is called severe preeclampsia if the systolic blood pressure is ≥ 160 mmHg and diastolic blood pressure ≥ 110 mmHg

accompanied by proteinuria of more than 5g / 24 hours (Peres, Mariana and Cairrão, 2018).

Eclampsia is an advanced condition of severe preeclampsia that is not resolved properly, leading to seizures. About 75% of eclamptic seizures occur before delivery, 50% within the first 48 hours after delivery, and can occur after 6 weeks postpartum. Eclampsia can lead to coma and even death, either before or after childbirth (Bernstein et al., 2017). The rate of caesarean sections in cases of severe preeclampsia is estimated to be high (around 70% or more in preterm pregnancies), because many obstetricians prefer performing caesareans in women with preeclampsia, even with a reassuring fetal status (Ferreira et al., 2016). Postpartum patients with severe preeclampsia-eclampsia are most at risk of infection due to childbirth with caesarean section, at risk of bleeding due to decreased ability of uterine



contractions due to administration of MgSO₄ and at risk of postpartum depression due to infant loss and inability to adapt to hormonal changes (Frawley et al., 2020; Hoedjes et al., 2011; Peres et al., 2018; Stern et al., 2014).

Several studies on the experiences of patients who received emergency treatment for severe preeclampsia indicated that patients felt that their lives were threatened, felt discomfort, tried to reduce discomfort, worried about the failure of treatment measures, felt close to death, felt that they did not get full support, expected friendly service and yearned for better conditions. The sense of loss can cause emotional distress to the mother, and leave a deep sense of grief. It is found that the patient expresses the need for professional support or social support (Frawley et al., 2020). To minimize the impacts, the needs and self-care of postpartum patients with severe preeclampsia and eclampsia should be identified as early as possible (Frawley et al., 2020). Assessment of the need and self-care of postpartum patients with severe preeclampsia and eclampsia is essential for professionals to meet the need and self-care of patients. Thus, an appropriate assessment instrument based on need and self-care of postpartum patients is needed. Based on a preliminary study on obstetric wards in several hospitals in Surabaya, there was no specific assessment instrument for postpartum patients with severe preeclampsia and eclampsia.

Wiedenbach's "need for help" model reveals that nurses are people who can help postpartum clients with preeclampsia-eclampsia overcome problems and improve their welfare through their actions, thoughts, feelings, words, writing, and body movements, while postpartum clients with preeclampsia-eclampsia are recipients of assistance (need for help) from health professionals either in the form of care, advice or education. Three goals put forward by Wiedenbach in helping clients are: (1) preventing maternal emergency, (2) reducing anxiety due to loss, (3) building effective coping in dealing with emergencies and losses by collaborating and coordinating with other health teams so that clients get actioned appropriate treatment procedures. In addition, Orem's "self-care" model sees that every individual has the ability and potential to care for himself and achieve prosperity (Alligood, 2018). In this case, the modification of Wiedenbach's need for help and Orem's self-care models can allegedly be applied to overcome the physical and psychological crisis and emergency problems of postpartum patients due to preeclampsia and eclampsia and help postpartum patients and their families achieve independence in self-care and baby care according to their abilities (Rahayu, 2015).

The development of assessment instruments has been carried out in several previous studies. Based on international joint commission accreditation standards, there is only a general nursing assessment instrument,

not specifically for postpartum patients (Rachmania, Sunaringtyas and Widayati, 2019). The development of maternal assessment instrument has been done in Brazil which is tracing about the quality of life of the mother and her baby, not identifying about emergency care needs as an impact of preeclampsia and eclampsia (Zubaran et al., 2009). Another study was conducted in Turkey aimed to compare the various existing instruments and determine which instrument is the most suitable for assessing the antenatal and postpartum period for Turkish society (Çakşak et al., 2018). Similar studies found the evaluation of changes in postnatal care using the Parents' Postnatal Sense of Security (PPSS) instrument. The study evaluated two models of postnatal care using a questionnaire incorporating the PPSS instrument. The PPSS can be used by mothers and fathers in the postpartum period to identify the sense of security of parents in the postnatal period (Kvist and Persson, 2009). However, those studies have not linked with the assessment instrument in postpartum with preeclampsia-eclampsia.

To overcome the limitation and the weakness of the previous studies, the current study aimed to develop an assessment instrument for postpartum patients with severe preeclampsia-eclampsia based on Wiedenbach's need for help and Orem's self-care models according to research with R&D design. This study is expected to identify the need for help and self-care of such patients by nurses and midwives in the emergency and maintenance period. Thus the risk of morbidity and mortality due to severe preeclampsia and eclampsia can be reduced significantly.

Materials and Methods

Study Design

The study used a research and development (R&D) design, a method used to produce certain products and test the effectiveness of these products (Yoshikawa, 2012). The purpose of using this method was to develop an assessment instrument in postpartum patients with pre-eclampsia based on Wiedenbach's need for help and Orem's self-care models. The research was conducted in two stages, namely the development stage and the trial stage.

At the development stage, the study used a qualitative research design and was carried out in the obstetric ward and the Medical Record Section of Airlangga University Hospital, Surabaya. Data collection was carried out by interviewing and tracing the contents of postpartum patient documents with preeclampsia-eclampsia.

The trial stage used a quasi-experimental, and the respondents were 60 midwives who were in charge for postpartum patients with preeclampsia in the obstetric ward of Airlangga University Hospital, Surabaya, and Haji Hospital, Surabaya.

Respondent

At the development stage, the participants were 30 midwives and 30 postpartum patients with severe preeclampsia-eclampsia, as well as 100 medical records stored in Airlangga University Hospital, Surabaya.

In the trial stage, the respondents were 60 midwives and 120 postpartum patients with preeclampsia in the obstetric ward of Airlangga University Hospital and Haji Hospital. The respondents were selected using purposive sampling technique, which is a sample selection method based on the predetermined criteria: Midwives who have experience caring for postpartum patients at least 1 year, as well as postpartum patients who have preeclampsia-eclampsia and are not admitted to the intensive care unit.

Instrument

The data were collected using some instruments based on the two stages. In the development stage, the instruments used were observation sheets, interview guidelines, and documentation. In the trial stage, the instrument used was a questionnaire to test the effect of the instrument of Assessment of Postpartum Mothers with Severe Preeclampsia and Eclampsia based on the "Need for Help" and "Self-Care" models.

Data Collection

The development stage began with a search for the availability of assessment instruments used by health workers in conducting studies of postpartum patients with preeclampsia-eclampsia. At this stage, in-depth interviews and documentation studies were carried out on the use of the assessment format, which had been used for the past 1 year. In-depth interviews were conducted with 30 midwives to explore their opinion on the instrument for assessing postpartum patients with severe preeclampsia that has been used so far. The media used at this stage was an interview guide that had been prepared based on the "need for help" and "self-care" models. To trace the documentation that used so far, a checklist was used. The results of in-depth interviews showed that the instruments provided by the hospital were general for all obstetric patients, not specific for postpartum patients with preeclampsia-eclampsia, so that often the risk of emergency due to preeclampsia was not detected.

The results of the qualitative analysis in the first stage were used as a basis for developing the new assessment instruments for postpartum patients with preeclampsia based on Wiedenbach's need for help and Orem's self-care models. After the new assessment instruments for the postpartum patients were arranged, a focus group discussion (FGD) was then carried out by involving researchers, patients, midwives, and experts on the preparation of the instruments. The results of the FGD were used to make improvements to the development of the assessment instrument.

At the second stage, it began with the midwife filling out a questionnaire about the old assessment instrument. Then the midwife was introduced to the new assessment instrument. After understanding about how to use the new assessment instrument, the midwife tried to conduct an assessment using the new instrument on two patients. After using the new assessment instrument, the midwife was asked to fill out a questionnaire about it.

Data Analysis

At the development stage, data were analyzed using qualitative method. The analysis consists of three activities that occur simultaneously: data reduction, data presentation, data withdrawal conclusion/verification. In this case, it was carried out to obtain conclusions about the use of assessment instruments that have been used so far. At the trial stage, data were analyzed using the Wilcoxon test with the significance level < 0.001 , because the result of the normality test of the post-test was not normal (Sugiyono, 2019). In this research, the validity test used the validity of the construction (construct validity) by asking for opinions from experts (experts' judgments). Instruments were made according to the aspects to be measured which were then consulted with experts to determine an opinion whether the instrument was feasible or not. In addition, testing the validity of the instrument was carried out using the Product Moment Correlation technique. If the results of $r_{count} > r_{table}$ have a significance level 5% it will be declared valid. In this study, the reliability of the instruments that have been compiled was tested by using the Cronbach's alpha technique. If coefficient reliability (r_{11}) is 0.7, then the instrument is declared reliable (Sugiyono, 2019).

Ethical Consideration

Prior to conducting the research, the research proposal was reviewed and received ethical approval from the Health Research Ethics Commission Team at Airlangga University Hospital No. 136/KEH/2019 and Haji Hospital Surabaya No. 073/34/KOM.ETIK/2019.

Results

The results of the development stage are described in [Table 1](#), [Table 2](#), and [Table 3](#), while the trial stage can be seen in [Table 4](#), [Table 5](#), and [Table 6](#).

The results of the evaluation of the old assessment instruments in the obstetric ward are shown in [Table 1](#). The finding showed that 5 of the 10 essential assessment components based on Wiedenbach's need for help and Orem's self-care models were not found. The old assessment instrument is (1) not specific assessment, (2) does not identify the risk of bleeding as an effect of administration of magnesium sulfate, (3) does not identify the ability to meet basic needs, (4) does not identify self-care abilities, and (5) does not identify parental readiness. Four components were partially

Table I Evaluation of old instrument

Instrument Standards	Information	Category
There is a focus or specific assessment format for postpartum patients with severe preeclampsia/eclampsia	The assessment format is not specific for postpartum patients with severe preeclampsia/eclampsia, still mixed with other cases (antepartum, intrapartum, and gynecology)	Not appropriate
Based on Wiedenbach's need for help model, the assessment of postpartum patients with severe preeclampsia/eclampsia should be sufficient to detect signs of eclampsia impending: <ol style="list-style-type: none"> 1. Headache complaint 2. Complaints of epigastric pain 3. Complaints of vision impairment 4. Complaints of nausea, vomiting 5. Help needed to deal with impending eclampsia in the hospital and after being discharged 	In the assessment format there is already a complaint assessment of dizziness and nausea and vomiting, but it is not specific for signs of impending eclampsia	Partially suitable
Based on Wiedenbach's need for help model, the assessment of postpartum patients with severe preeclampsia/eclampsia should be sufficient to identify fluid balance: <ol style="list-style-type: none"> 1. Intake 2. Output 3. Assistance needed for fluid balance in the hospital and after hospital discharge 	In the assessment format, there is already an assessment of fluid intake and output	Suitable
Based on Wiedenbach's need for help model, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify signs of magnesium sulfate therapy poisoning: <ol style="list-style-type: none"> 1. Respiratory depression (RR <12 times/minute) 2. Tendon reflexes disappear 3. Declining awareness 4. Help needed to overcome magnesium sulfate intoxication 	In the assessment, there has been an assessment of breathing, awareness, but there is no assessment of tendon reflexes	Partially suitable
Based on Wiedenbach's need for help model, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify the risk of bleeding as an effect of the administration of magnesium sulfate <ol style="list-style-type: none"> 1. Uterine contractions 2. Height of the uterine fundus 3. Characteristics of vaginal discharge 4. Assistance needed to deal with the risk of bleeding in the hospital and after being discharged 	In the assessment format, there is no assessment of uterine contractions, the height of the uterine fundus, or characteristics of vaginal discharge for postpartum	Not appropriate
Based on Wiedenbach's need for help model, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify the risk of pulmonary edema as a hypoalbumin effect <ol style="list-style-type: none"> 1. Fluid restriction 2. Lung sounds 3. Respiration rate (RR) 4. Assistance needed to deal with the risk of pulmonary edema in the hospital and after hospital discharge 	In the assessment, there is an assessment of lung sounds and RR, but there is no assessment of fluid restriction	Partially suitable
Based on Wiedenbach's need for help model, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify the ability to meet basic needs <ol style="list-style-type: none"> 1. Oxygenation 2. Nutrition 3. Elimination 4. Mobilization 5. Rest and sleep 6. Assistance needed to meet basic needs in the hospital and after being discharged 	In the assessment, there is an assessment of nutrition, but there is no assessment of oxygenation, elimination, mobilization, health, and sleep, and the assistance needed	Not appropriate

Table 1 Evaluation of old instrument (continuos)

Instrument Standards	Information	Category
Based on Wiedenbach's need for help model, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify comfort 1. Pain scale 2. Causes of pain 3. The ability and habit of dealing with pain 4. Assistance needed for comfort in the hospital and after being discharged	In the assessment, there is an assessment of the scale of pain but there is no assessment of the causes of pain, as well as the ability and habits to deal with the pain	Partially suitable
Based on Wiedenbach's need for help and Orem's self-care models, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify abilities to meet basic care 1. Personal hygiene 2. Perineal care 3. Wound care 4. Knowledge of postpartum danger signs 5. Help is needed about self-care in the hospital and after being discharged	In the study, there was no assessment of personal hygiene, milk production and breastfeeding, and perineal care/wound care	Not appropriate
Based on Wiedenbach's need for help and Orem's self-care models, the assessment of postpartum patients with preeclampsia/eclampsia should be sufficient to identify parental readiness 1. Mental readiness 2. Readiness of knowledge about baby's needs and care 3. Breast Production and Breastfeeding 4. Support system 5. Rocks needed for baby care in the hospital and after hospital discharge	In the assessment format, there is no assessment of parental readiness	Not appropriate

Table 2 First focus group discussion results

Strategic Issues	Causes	FGD results	Researcher Study
Specific assessment instrument for postpartum with severe preeclampsia-eclampsia	1. The unavailability of a special assessment instrument for postpartum patients with severe preeclampsia/eclampsia 2. The instrument of the outpatient assessment is made public for all cases treated in the midwifery ward	Development of specific assessment instruments for postpartum patients with severe preeclampsia/eclampsia	Development of a special assessment instrument is needed to identify the needs, abilities, independence, and assistance needed by postpartum patients with severe preeclampsia -eclampsia.
Self-management support	There are no instruments that show the self-management data of postpartum patients with severe preeclampsia/eclampsia who need support	Development of an instrument to explore the self-management data of postpartum patients with severe preeclampsia/eclampsia who need support	Development of the instrument needs to be done in the obstetric ward to overcome the emergency conditions and improve the ability and independence of postpartum patients with severe preeclampsia/eclampsia
Decision support	No instrument shows the decision support in the form of a written list of things that must be studied about the needs, abilities, and assistance needed by postpartum patients with severe preeclampsia/eclampsia	Development of Instrument which includes: 1. List of patient needs 2. List of patient abilities 3. List of assistance the patient needs	Development of an Instrument is needed to overcome the emergency conditions and improve the ability and independence of postpartum patients with severe preeclampsia/eclampsia
Service system design	No instrument shows the things that need to be studied about the needs, abilities, and assistance needed by postpartum patients with severe preeclampsia/eclampsia	1. Development of Instrument that includes a list of patient needs 2. List of patient abilities 3. List of assistance the patient needs	Development of an Instrument is needed to overcome the emergency conditions and improve the ability and independence of postpartum patients with severe preeclampsia/eclampsia
Clinical information	Clinical information such as signs of eclampsia impending, self-care ability, parental readiness has not been systematically arranged	Systematic development of clinical information 1. Subjective data 2. Objective data	Development of Instrument is carried out to determine the patient's development and arrange further self-management strategies

Table 3 Results of expert discussions

Instrument	Component	Feedback
Specific or focus assessment instrument for postpartum patients with severe preeclampsia-eclampsia	1. Self-management support	Specific data based on the model need for help and self-care
	2. Decision support	Data that examines the management of postpartum patients with severe preeclampsia/eclampsia which includes: <ol style="list-style-type: none"> 1. Symptoms of impending eclampsia 2. Fluid balance 3. Signs of magnesium sulfate poisoning 4. Risk of pulmonary edema 5. Risk of bleeding 6. Self-care ability 7. Comfort 8. Parental readiness 9. Meeting basic needs
	3. Service system design	Development of instruments is needed to improve the alarm warning to detect early and overcome emergencies and improve the ability of postpartum patients with severe preeclampsia/eclampsia in the obstetric ward
	4. Clinical information	An instrument is available that examines data about: <ol style="list-style-type: none"> 1. Symptoms of impending eclampsia 2. Fluid balance 3. Signs of magnesium sulfate poisoning 4. Risk of pulmonary edema 5. Risk of bleeding 6. Self-care ability 7. Comfort 8. Parental readiness 9. Meeting basic needs

Table 4 Normality test results

	Pre	Post
Kolmogorov-Smirnov Z	1.133	1.701
Asymp. Sig. (2-tailed)	0.153	0.006
Information	Normal	Abnormal

Table 5 Statistical test results

Group	N	Mean Rank	Z score	P-value
Old Instrument	100	19.93	-5.379	<0.001
New Instrument	100	29.72		

Table 6 Second FGD results

Strategic Issues	Cause	FGD results	Researcher
Specific assessment instrument or focus for postpartum patients with severe preeclampsia/eclampsia	It cannot be applied directly to postpartum patients with severe preeclampsia/eclampsia	It has been done even though it has not been a supplementary document	The assessment instrument of the postpartum patient with severe preeclampsia/eclampsia based on the "Need For Help" and "Self-Care" Models can already be used in the obstetric ward to get complete information from the patient

compliant, namely (1) detecting signs of impending eclampsia, (2) signs of magnesium sulfate therapy poisoning, (3) the risk of pulmonary edema as a hypoalbumin effect, and (4) comfort. There is only one component which was suitable, namely about fluid balance. Therefore, the old instrument is insufficient to assess the need for help and self-care of postpartum patients with severe preeclampsia/eclampsia.

The results of the FGD in the first stage are shown in [Table 2](#). The finding showed that, based on five aspects of strategic issues, the development of an instrument to assess postpartum patients with severe preeclampsia or eclampsia was needed. The FGD recommendations are the approval of the development of a specific assessment instrument for postpartum patients with severe preeclampsia/eclampsia. Furthermore, a new assessment instrument was developed for postpartum patients with severe-eclamptic preeclampsia based on Wiedenbach's need for help and Orem's self-care models.

The results of discussions with experts on the four elements are shown in [Table 3](#). The finding showed that the development of assessment instruments is needed to improve the alarm warning to detect early and overcome emergencies and improve the ability of postpartum patients with severe preeclampsia/eclampsia in the obstetric ward. Recommendations from an expert discussion on the instrument for the study of postpartum patients with severe preeclampsia/eclampsia were: 1) the development of a specific assessment tool for postpartum patients with severe preeclampsia/eclampsia, and 2) the components of the instrument should include: self-management support, decision support, service system design, and clinical information.

Before further analysis was carried out, the normality of the data was tested with the Kolmogorov Smirnov test because the data are more than 50. The data normality test can be seen at [Table 4](#) and the test result at [Table 5](#). In this study, the results of the pre-test data normality test were normally distributed, while the post-test was not normal. So that the statistical test used is the Wilcoxon Signed Rank Test. The old assessment instrument was rated lower with a mean rank of 19.93, while the new assessment instrument was with a mean rank value of 29.72. The Wilcoxon test results obtained a value of -5.379 with $p < 0.001$. So there is a significant difference between the old assessment instrument and the new postpartum patient assessment instrument.

The results of the second stage FGD are shown in [Table 6](#). The finding shows the assessment instrument can be used in obstetric wards to get the more complete information from the patients. In addition, the experts recommended to improve the instruments by adding data components about the medication that patients should continue at home. This is important because treatment with antihypertensive drugs is recommended for severe postpartum hypertension.

Discussions

The results of this study illustrate that an instrument carried out to assess postpartum patients with severe preeclampsia/eclampsia was inadequate. Some previous studies have described that postpartum patients with preeclampsia are at risk for experiencing emergencies, namely eclampsia, bleeding, and dyspnea (Cairns et al., [2017](#); NICE, [2019](#); Kvist & Persson, [2009](#); Magee et al., [2014](#); Rana et al., [2019](#); Rawlins et al., [2018](#)).

Based on the results of previous research, it illustrates that the need for help and self-care models are effective to be applied to patients with maternal complications, both physical and psychological. This model can be used in patients with preeclampsia, bleeding, preterm labor, and in patients with postpartum depression (Machmudah, [2015](#); Rahayu, [2015](#); Wulandari and Setyowati, [2015](#); Dwi Susanti et al., [2016](#)).

The results of this study illustrate that the assessment instruments that have been used so far are still general and not entirely specific to identifying the special needs of preeclamptic patients. Based on the input from the experts, the development of instruments for postpartum patients with severe preeclampsia/eclampsia using the "Need for Help" concept proposed by Wiedenbach with the goals in helping clients are: 1) to prevent maternal emergencies, 2) reduce anxiety due to loss, and 3) develop effective coping by collaborating and coordinating with other health teams so clients can get appropriate action and handling procedures.

In addition, the self-care theory by Orem, which believes that every individual has the ability and potential to care for themselves and achieve prosperity, was also used to develop the instruments. Changes that occur during the postpartum period cause a decrease in the client's independence to meet their needs. Nurses play a role in helping to increase the independence of clients to meet their self-care needs through a learning process or exercise in the form of self-care, creating an environment that facilitates the achievement of independence.

The statistical test showed that there is a significant difference between the old assessment instrument and the new one, which was considered more effective. Based on the results of data analysis from the results of filling out the questionnaire, almost all respondents gave the opinion that the assessment instrument for postpartum patients with severe preeclampsia-eclampsia based on Wiedenbach's need for help and Orem's self-care models met the criteria of functionality, efficiency, and usability. The effectiveness of using the need for help and self-care models in providing nursing care has been illustrated in the results of previous studies in different cases (Çapik et al., [2015](#); Ferreira et al., [2016](#); Hajira Irshad Ali, [2018](#)).

Conclusions

The assessment instruments in obstetric ward have not been specific for postpartum patients with severe

preeclampsia/eclampsia, while the new instrument has been developed for postpartum patients with severe preeclampsia-eclampsia based on Wiedenbach's need for help and Orem's self-care models through FGD and expert discussion. The new assessment instruments met the criteria of functionality, efficiency, and usability.

For stakeholders, midwives recommend the use of the assessment instrument of the postpartum patients with severe preeclampsia-eclampsia based on Wiedenbach's need for help and Orem's self-care models in the obstetric ward setting.

Acknowledgment

Thanks are conveyed to the nurses/midwives in the obstetrics ward and management, as well as postpartum patients both at the Airlangga University Hospital and the Haji Hospital Surabaya.

Funding Source

The research was provided financial support by Poltekkes Kemenkes Surabaya.

References

- Alligood, M. R. (2018) *Nursing Theorists and Their Work*. 9th edn. St. Louis, MO: Elsevier Inc.
- Cairns, A. E. *et al.* (2017) 'Postpartum management of hypertensive disorders of pregnancy: A systematic review', *BMJ Open*. BMJ Publishing Group. doi: 10.1136/bmjopen-2017-018696.
- Çakşak, A. *et al.* (2018) 'Tools for assessing parenting in the prenatal and postpartum period in Turkey', *New Trends and Issues Proceedings on Advances in Pure and Applied Sciences*, (10), pp. 104–109. doi: 10.18844/gjpaas.v0i10.3750.
- Çapik, A. *et al.* (2015) *The Effect of the Care Given Using Orem's Self-Care Model on the Postpartum Self-Evaluation*, *International Journal of Caring Sciences*.
- Dwi Susanti, H. *et al.* (2016) 'Wiedenbach Need for Help and Swanson Caring Model on Nursing Care of Prolonged Second Stage of Labour', 7(1).
- Ferreira, M. B. G. *et al.* (2016) 'Nursing care for women with pre-eclampsia and/or eclampsia: Integrative review', *Revista da Escola de Enfermagem*. Escola de Enfermagem de Universidade de Sao Paulo, pp. 320–330. doi: 10.1590/S0080-623420160000200020.
- Frawley, N., East, C. and Shaun Brennecke (2020) 'Women's experiences of preeclampsia: a prospective survey of preeclamptic women at a single tertiary centre', *J Obstet Gynaecol*, 20(1), pp. 65–69. doi: 10.1080/01443615.2019.1615040.
- Hajira Irshad Ali, B. (2018) 'Application of Orem Self Care Deficit Theory on Psychiatric Patient', *Ann Nurs Pract*, 5(1), p. 1093.
- Health Ministry (2021) 'Profil Kesehatan Propinsi Jawa Timur 2019', date accessed 25 May 2021.
- Hoedjes, M. *et al.* (2011) 'Postpartum Depression after mild and severe preeclampsia', *Journal of Women's Health*, 20(10), pp. 1535–1542. doi: 10.1089/jwh.2010.2584.
- Kvist, L. J. and Persson, E. K. (2009) 'BMC Pregnancy and Childbirth Evaluation of changes in postnatal care using the " Parents ' Postnatal Sense of Security " instrument and an assessment of the instrument ' s reliability and validity', 10, pp. 1–10. doi: 10.1186/1471-2393-9-35.
- Machmudah (2015) 'Penerapan Model Konsep Need for Help Dan Self Care Pada Asuhan Keperawatan Ibu Pre Eklampsia Berat Dengan Terminasi Kehamilan', *Jurnal Keperawatan Maternitas*, 3(1), pp. 16–26.
- Magee, L. A. *et al.* (2014) 'Diagnosis, Evaluation, and Management of the Hypertensive Disorders of Pregnancy: Executive Summary', *Journal of Obstetrics and Gynaecology Canada*. Elsevier Inc, 36(5), pp. 416–438. doi: 10.1016/S1701-2163(15)30588-0.
- NICE. *Hypertension in pregnancy: diagnosis and management NICE guideline* (2019).
- Peres, G., Mariana, M. and Cairrão, E. (2018) 'Pre-Eclampsia and Eclampsia: An Update on the Pharmacological Treatment Applied in Portugal', *Journal of Cardiovascular Development and Disease*. MDPI AG, 5(1), p. 3. doi: 10.3390/jcdd5010003.
- Peter S. Bernstein, James N. Martin, John R. Barton, Laurence E. Shields, Maurice L. Druzin, Barbara M. Scavone, Jennifer Frost, Christine H. Morton, Catherine Ruhl, Joan Slager, Eleni Z. Tsigas, Sara Jaffer, M. K. M. (2017) 'Consensus Bundle on Severe Hypertension During Pregnancy and the Postpartum Period', *Journal JOGNN - Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 46(5), pp. 776–787. doi: 10.1016/j.jogn.2017.05.003.
- Rachmania, D., Sunaringtyas, W. and Widayati, D. (2019) 'Pengembangan instrumen pengkajian M3 (METHOD) dalam manajemen keperawatan berbasis standar akreditasi joint comission international', *The Indonesian Journal of Health Science*, 11(2), pp. 143–154.
- Rahayu, K. D. (2015) 'Theory implementation of need for help wiedenbach and self care orem in pregnancy case with tuberculosis multi drug resistance', *JMCRH*, 1(118–129).
- Rana, S. *et al.* (2019) 'Preeclampsia: Pathophysiology, Challenges, and Perspectives', *Circulation Research*. Lippincott Williams and Wilkins, 124(7), pp. 1094–1112. doi: 10.1161/CIRCRESAHA.118.313276.
- Rawlins, B. *et al.* (2018) 'Screening and management of pre-eclampsia and eclampsia in antenatal and labor and delivery services: Findings from cross-sectional observation studies in six sub-Saharan African countries', *BMC Pregnancy and Childbirth*. BioMed Central Ltd., 18(1). doi: 10.1186/s12884-018-1972-1.
- Shamsi, U., Saleem, S. and Nishter, N. (2013) 'Epidemiology and Risk Factors of Preeclampsia ; An Overview of Observational Studies', *Al Ameen J Med Sci*, 6(4), pp. 292–300.
- Stern, C. *et al.* (2014) 'The impact of severe preeclampsia on maternal quality of life', *Quality of Life Research*. Kluwer Academic Publishers, 23(3), pp. 1019–1026. doi: 10.1007/s11136-013-0525-3.
- Sugiyono (2019) *Metodologi Penelitian Kuantitatif, Kualitatif, dan R&D*. 1st edn. Bandung: ALVABETA.
- Wulandari, P. and Setyowati (2015) 'Penerapan model keperawatan', *Versi online / URL: http://ejournal.umm.ac.id/index.php/keperawatan/article/view/2346*, 5(2), pp. 173–177.
- Yoshikawa, H. (2012) *Design Methodology for Research and Development Strategy — Realising a Sustainable Society —*.
- Zubaran, C. *et al.* (2009) 'An assessment of maternal quality of life in the postpartum period in southern Brazil: A comparison of two questionnaires', *Clinics*, 64(8), pp. 751–756. doi: 10.1590/S1807-59322009000800007.

How to cite this article: Suprihatin, E., Wuryaningsih, S. R. (2022) 'Development assessment instrument postpartum patients with severe preeclampsia-eclampsia based on need for help and self-care models', *Jurnal Ners*, 17(1), pp. 47–54. doi: <http://dx.doi.org/10.20473/jn.v17i1.35029>

Factors associated with compliance in implementing health protocols to prevent COVID-19 in Indonesia: A cross-sectional study

Ambar Dwi Erawati ^{1,*} 

¹ Widya Husada Semarang University, Indonesia

*Correspondence: Ambar Dwi Erawati. Address: Subali Raya Street No.12, West Semarang 50146, Indonesia.

Email: ambarerawati@gmail.com

Responsible Editor: Ilya Krisnana

Received: 23 February 2021 ○ Revised: 16 April 2022 ○ Accepted: 18 April 2022

ABSTRACT

Introduction: To prevent the COVID-19 spread, the government requires people to comply with health protocols in public facilities, namely by wearing masks, washing hands and keeping a distance. This study aimed to determine what factors affect and reason for compliance in implementing health protocols.

Methods: This research was an analytical survey research with a cross-sectional design conducted in Semarang City, Indonesia, with a total sample of 400 respondents. The independent variables were age, education, occupation, and health status. The dependent variable was compliance in wearing masks, washing hands and keeping a distance. Data were analysed with logistic regression test with significance level of 0.05.

Results: From the survey, 76.6% of respondents obeyed wearing masks, 57.4% of respondents obeyed to wash their hands and 71.1% of respondents obeyed to keep their distance. The backward type logistic regression test found that the factors associated with health protocol compliance were age, education, occupation, health status, existing regulations and the desire to maintain health with a significance value of <0.001. However, field of work was not associated with using masks ($p = 0.273$), while health status was not associated with handwashing habits ($p = 0.344$).

Conclusions: Factors that are associated with respondents' compliance with the health protocol are age, education, occupation, health status, self-motivation and regulation. The government need to raise awareness on adhering health protocols through the promotion about healthy lifestyle.

Keywords: adherence; COVID-19; health awareness; health protocols

Introduction

COVID-19 will not only have an impact on the health sector, but will also have an impact on the social, economic, educational and other fields (Ministry of Fianance, 2020). Research conducted by Sardar, Ray, Hasan, and Chitra (2022) shows that there is a reduction in the number of employees at companies even though online sales have increased. The government has formulated a strategy by making policies to deal with COVID-19 in the field of information and communication with a special task force to COVID-19 so that it can be accepted by the wider community by broadcasting live on TV, the government's official website, and creating an Android-based application for detecting COVID-19 events. Mass media have reported a lot of discussion

about the COVID-19, including the number of COVID-19 cases, prevention, health protocols, and vaccinations.

In order to break the chain and deal with COVID-19, the government through the task force prepares health protocols for every activity that potentially transmit the disease. For the prevention of COVID-19 in public places and public facilities, the government has prepared a health protocol for the community in public places and facilities (Ministry of Health, 2020). By implementing the health protocol, it is expected the people in a country to be able to avoid COVID-19 infection, for which the health protocol must be adhered. However, people's compliance towards the COVID-19 health protocol has not been widely studied.



Compliance to implement health protocols is part of community behavior in preventing COVID-19. A study found that knowledge affects behavior (Istiqomah and Notobroto, 2016) and there is a significant positive relationship between education level and healthy living behavior (Putri, 2017). Knowledge has significantly associated with the behavior of using personal protective equipment (PPE). Based on the results of this study, knowledge plays an important role in people's behavior, knowledge influenced by information obtained from the community (Erawati, Alfiani and Kurniasih, 2020). However, it is different from previous research that good knowledge is not in line with good behavior (Erawati, Puspita and Cahyaningsih, 2020).

Based on this background, the aim of this study was to analyse what factors and reasons were associated with

adherence in implementing health protocols to prevent COVID-19.

Materials and Methods

Study Design

This study was a quantitative research with an analytical survey approach with a cross-sectional design. The independent variables were age, education, occupation and health status. The dependent variable was compliance with the use of masks, hand washing and keeping a distance followed by reasons for compliance.

Table 1 Distribution of age, education, occupation, and health conditions (N = 400)

Variable	n	%
Age		
20-25 years	18	4.5
26 - 35 years	76	19
36 - 45 years	204	51.1
46 - 55 years	51	12.7
56 - 59 years	51	12.7
Education level		
SMP (Junior High School) or lower	34	8.5
SMA (Senior High School)	68	17.1
Universities	298	74.4
Occupation		
Public Services (security, banking)	60	14.9
Industry	17	4.3
Entrepreneur (trade, etc.)	25	6.4
Education	128	31.9
Health	34	8.5
Others (including Housewives)	136	34.0
Health Status		
Suffering from a disease	44	8.5
Healthy	366	91.5

Table 2 Distribution of health protocol adherence (N = 400)

Health Protocol Adherence	n	%
Wearing a Mask		
Always	306	76.6
Often	34	8.5
Sometimes	52	12.8
Never	8	2.1
Washing Hands		
Always	230	57.4
Often	68	17.0
Sometimes	94	23.4
Never	8	2.2
Keeping a Distance		
Always	284	71.1
Often	44	10.9
Sometimes	55	13.7
Never	17	4.3

Table 3 Distribution of reasons for complying with health protocols (N = 400)

Health Protocol	Reason			
	Regulations		Health intention	
	n	%	n	%
Wearing a Mask	56	14.9	340	85.1
Washing Hands in Public Facilities	56	14.9	340	85.1
Keeping a Distance in Public Facilities	68	17.0	332	83.0

Table 4 The results of data analysis of age, education, occupation and health status on the health protocol adherence

Variable	B	t	p-value
Wearing Mask			
Age	0.853	29.403	<0.001
Education	-0.648	-16.986	<0.001
Occupation	0.472	6.361	<0.001
Field of work	0.023	1.097	0.273
Health status	0.172	6.456	<0.001
Washing Hands			
Age	0.514	12.961	<0.001
Education	-1.054	-21.071	<0.001
Occupation	0.460	19.996	<0.001
Field of work	0.464	19.887	<0.001
Health status	0.078	0.948	0.344
Keeping a Distance			
Age	0.767	18.415	<0.001
Education	-0.858	-16.342	<0.001
Occupation	0.171	6.982	<0.001
Health status	0.558	6.486	<0.001

Table 5 Test results analysis of reasons for the habit of using masks, washing hands and keeping a distance

Variable	B	t	p-value
Hand wash	3.661	13.790	<0.001
Regulations	0.537	5.718	<0.001
Health	-1.599	-17.404	<0.001
Using Mask	4.892	37.122	<0.001
Regulations	0.126	2.710	0.007
Health	-2.009	-44.054	<0.001
Keeping a Distance	4.922	32.773	<0.001
Regulations	0.190	3.616	<0.001
Health	-2.056	-39.333	<0.001

Respondents

The study was conducted in the city of Semarang, with a population of 1,096,637 residents. After calculating with the Slovin's formula, the number of samples was 400 residents by accidental sampling (non-randomized). Inclusion criteria in sampling are residents of Semarang City aged 20-59 years who are productive. The exclusion criteria in this study were residents of Semarang City aged 20-59 years who were unproductive, getting sick, or not working.

Instrument

The research instrument used a closed questionnaire which was distributed online using a Google Form ([Supporting Information 1](#)). The form contains the first part, namely Informed Consent and the demographic characteristics of the respondents (age, occupation, education and health status). The second part consists of the habit of using masks, washing hands and keeping a distance. All independent variables are classified in [Table 1](#). Questions related to compliance with using masks, washing hands and keeping a distance are classified as always (4), often (3), sometimes (2), and never (1), followed by the reason for compliance (i.e. regulation or health reasons).

The validity of the previous questionnaire was tested in 30 respondents from 400 respondents as sample of this study with the help of Pearson's SPSS Product

Moment, it was obtained that arithmetic mean was 0.369 -0.976 or greater than r table, which is 0.349 (valid). Reliability test was by looking at the value of Cronbach's alpha and obtained a value of 0.897 (reliable).

Data Collection

Data collection was carried out within two months (November-December 2020). Data retrieval by accidental sampling was distributed in several WhatsApp groups with the help of 15 enumerators who had different characteristics (age and occupation) to avoid data bias in certain groups. The 15 enumerators were previously known to the researcher. The researcher coordinated with the enumerators regarding the explanation of the questionnaire, the enumerator's rights and the tasks carried out by the enumerator by distributing the questionnaire link via WhatsApp.

Data Analysis

The data analysis carried out was univariate and bivariate analysis. The distribution of characteristics, distribution of adherence and distribution of reasons for complying with the protocol were carried out by univariate analysis. To see what factors associated with the health protocol, bivariate analysis was carried out by performing a backward type logistic regression test with the help of SPSS 16. The t-table value with a sample of 400 was 1.966, and the researcher used a value of $\alpha = 0.05$. so that, if the analysis results get a t value of more than 1.966

or less than 0.05, it will be used as a factor that affects compliance in implementing health protocols. If the t value is less than 1.966 or is more than 0.05, it is not used as a factor that is associated with compliance in implementing health protocols.

Ethical Consideration

This research has passed the ethical test with no. 60/EC-LPPM/UWHS/XI-2020 was issued by the Ethics Committee of Widya Husada University Semarang.

Results

[Table 1](#) shows the age characteristics of the majority of respondents are 36-45 years old, namely 51%. The majority of respondents were highly educated (74.4%). Regarding job characteristics, the majority of the respondents worked outside the field of education or internal industry, namely 34%. The characteristics of the health conditions of the majority of respondents are healthy, namely 91.5%.

Based on [Table 2](#), it is known that the majority of respondents comply with health protocols, both in using masks, washing hands and maintaining distance; 76.6% of respondents always use masks, 57.4% wash their hands in public places and 71.1% of respondents keep their distance in public places.

[Table 3](#) shows the majority of respondents comply with health protocols because they want to maintain their health to avoid COVID-19, which is 85.1%, but there are still those who comply because of regulations, which are 14.9% on compliance with wearing masks and washing hands and 17% on adherence to social distancing.

With a sample of 400 respondents, it was found that the df was 393 and the t-value was 1.966. After carrying out a backward test between the use of masks and the variables of age, education, occupation and health status simultaneously, it was found that the field of work had a t-value of less than 1.966 and a significance value was obtained of 0.273 or greater than 0.05 so that the field of work was not a factor affecting the use of masks in the prevention of COVID-19. However, age, education and health status are factors associated with respondents in wearing masks for the prevention of COVID-19.

Backward type logistic regression test between the variables of age, education, occupation and health status on hand washing found that the health status variable had a t-value of 0.948 or less than 1.966 and had a significance value of 0.344 or more than the probability value, which was 0.05 so that health status was not a factor that associated with respondents' compliance in washing hands in public facilities to prevent COVID-19. However, age, education, field of work are factors that were associated with respondents' handwashing habits in public facilities in preventing COVID-19.

Backward test between the variables of age, education, occupation and health status on maintaining distance in public places found the t-value in all independent variables is more than 1.966 and the significance value in all variables is less than the probability value. So age, education, occupation and health status are factors that were associated with social distancing habits in public places.

After a regression test was conducted between the habit of using masks and the reasons for complying with the regulations, it gave a significance value of 0.07 or more than 0.05. The habit of using a mask with a desire to be healthy gives a significance value of <0.001. The habit of washing hands, the habit of keeping a distance from the reasons for complying with regulations and maintaining health gives a significance value of <0.001.

Discussions

Factors Associated with Mask Use

The majority of respondents implemented health protocols, with an average of 68.4%. However, the habit of washing hands occupies the lowest average, which is 57.4%. This means that from all respondents there are those who always use masks, always keep their distance but don't always wash their hands.

According to UNICEF, the spread of COVID-19 is through splashes of phlegm containing the virus that enter the body through the eyes, nose or throat. However, it most often occurs through hands contaminated with droplets that eventually enter the body through the eyes, nose or throat (UNICEF, [2020](#)). Droplets that are splashed directly can cause droplets to stick to objects which are then held by other people who are not affected by COVID-19. Droplets enter the oral or nasal mucosa or the conjunctiva of the eye. For this reason, the government made rules for all residents to always wear masks and wash their hands (Ministry of Health, [2020](#)).

Based on the results of the backward test, it was found that the job characteristics had a significant value of 0.273 or more than 0.05, which means that work does not have a significant effect on the habit of using masks. The characteristics associated with the habit of using masks in this study are the characteristics of age, education and health status. This supports the research conducted by Juanda ([2020](#)) that education has a positive association on compliance and supports Notoatmodjo's theory that education is an exogenous factor that influences behavior (Notoadmojo, [2012](#)). Research conducted by Ozdemir, Ng, and Chaudhry ([2020](#)) found that the strongest indicator of behavior change in preventing COVID-19 is that respondents who are older, highly educated, anxious and married report higher adoption/frequency of preventive measures.

Work does not affect the use of masks, meaning that masks have become a habit of the people during the

pandemic in 2020, Even though working in health, education, industry or other occupations does not have a significant effect on wearing masks. The reason for wearing masks is 85.1% for maintaining health, meaning that the community already understands the importance of masks in preventing COVID-19.

Factors Associated with Handwashing

The results of the backward test show that the characteristics of age, education and occupation have a significantly smaller value than the probability so that it can be interpreted that age, education and occupation have an effect on hand washing habits, but health status has a significance value greater than the probability value, which is 0.344. So that health status does not affect respondents in washing their hands.

Respondents who have poor health conditions do not care about washing their hands. Although people who have inherited diseases have a higher risk of getting worse if they get COVID-19 (Zandkarimi, 2020).

The results of this study are not in line with research conducted by Riyadi and Larasaty (2020) that health status has a positive association with health protocol compliance. If it is associated with 14.9% of respondents who obediently wash their hands for reasons of regulations, it is necessary to give rules for mandatory hand washing areas, because so far public places only the areas where masks are mandatory.

Factors Associated with Social Distancing

The results of the backward test showed that the characteristics of age, education, field of work and health status can predict respondents behaviour in complying social distancing. This supports previous studies that state health behavior is associated with age, education, occupation and health status. Supporting research conducted by Dewi, Adawiyah, and Rujito (2019) found age has a significant influence in complying with the use of PPE and research conducted by Mualama (2020) showed that the majority of adults in his research have a healthy lifestyle (Muamala, 2020). Recommendations from the WHO that maintaining physical distance, wearing face masks, avoiding crowds, and practicing hand hygiene are forms of controlling COVID-19, the correct use of masks is highly recommended (Claeson, 2021).

Reasons for Implementing Health Protocols

In this study, showing compliance in implementing health protocols because they want to maintain health or to avoid COVID-19 is higher than because of the regulation factor can mean that respondents are aware of the causes of COVID-19 transmission even though there are still those who carry out health protocols because of regulations. However, from the results of the regression

test, it was found that the factor complying with the regulations on the habit of wearing masks got $p = 0.07$. It means that respondents think they can maintain their health and prevent COVID-19, only by wearing masks.

Contrary to what Claeson and Hanson (2021) said, the involvement of policy makers has an effect on changing people's behavior and research conducted by Stewart et al. (2020) found the need for collaboration from all parties, both health workers, government, communities, to form learning experiences for the community. The government sets rules for all citizens to always wear masks and wash their hands (Ministry of Health, 2020).

Socialization is an important step in implementing policies. Government and public health efforts in encouraging the prevention of COVID-19 are carried out by providing counseling or socialization (Sriarumtias et al., 2020). Social media is one of the factors facing COVID-19 (Addis and Abate, 2021).

The socialization of COVID-19 prevention by always wearing masks for respondents has been successful; however, prevention of always washing hands and keeping a distance in public facilities still needs to be improved because there are still respondents who obey hand washing and keep their distance for reasons of only complying with regulations.

This study did not examine implementation policies in the prevention of COVID-19 so it cannot know what can be known in compliance with implementing health protocols.

Conclusion

Age, education, occupation and health status are compliance factors in implementing health protocols to maintain distance, but field of work was not associated with using masks, while health status was not associated with compliance in washing hands. People must be aware that maintaining health begins with self-prevention, not only because of obeying the rules. This study suggest that the government need to encourage and raise awareness about personal health than regulation enforcement. The socialization of preventing COVID-19 by always wearing masks has been successfully carried out, but not yet on the habit of washing hands and keeping a distance.

Supporting Information

[Supporting Information 1](#)

References

- Addis, Y. and Abate, D. (2021) 'Social Work Responses and Household-level Determinants of Coronavirus Preparedness in Rural Ethiopia.', *Social work in public health*. Routledge, 00(00), pp. 1–13. doi: 10.1080/19371918.2021.1881014.
- Dewi, I. P., Adawiyah, W. R., & Rujito, L. (2019). Analysis of Adherence Level of Using Personal Protective Equipment for Dentist Profession Students at Unsoed Dental and Oral Hospital. *JEconomic Journal for Business and Accountants (JEBA)*, 21(4), 1–10. Retrieved from <http://jp.feb.unsoed.ac.id/index.php/jeba/article/viewFile/1541/1550>

- Erawati, A. D., Alfiani, N. and Kurniasih, D. (2020) 'Pengetahuan Ibu Hamil tentang Buku Kesehatan Ibu dan Anak', *Jurnal Kesehatan Masyarakat*, 19(02), pp. 50–54. doi: <https://doi.org/10.33221/jikes.v19i02.474>.
- Erawati, A. D., Puspita, D. and Cahyaningsih, O. (2020) 'Pengetahuan nifas tentang perawatan tali pusat di Puskesmas Ambarawa Kabupaten Semarang', *The Public Health Science Journal*, 9(01), pp. 43–47. doi: <https://doi.org/10.33221/jikm.v9i01.476>.
- Ministry of Finance (2020) 'Pernyataan Menteri Keuangan'. Ministry of Finance.
- Ministry of Health (2020) *Pedoman Pencegahan dan Pengendalian Corona Virus Disese (COVID-19)*, Ministry of Health.
- Istiqomah, N. and Notobroto, H. B. (2016) 'Pengaruh Pengetahuan, Kontrol Diri terhadap Perilaku Seksual Pranikah di Kalangan Remaja SMK di Surabaya', *Jurnal Biometrika dan Kependudukan*, 5(2), p. 125. doi: <http://dx.doi.org/10.20473/jbk.v5i2.2016.125-134>.
- Juanda, S. F. E. (2020) 'Factors that Influence the Adherence of Pregnant Women in Consuming Iron Supplements : Systematic Literature Review', *PLACENTUM health scientific journals and their applications*, 8(1), pp. 18–24.
- Mariam Claeson, S. H. (2021) 'COVID-19 and the Swedish enigma', *The Lancet*, 397(10271), pp. 259–261. doi: 10.1016/S0140-6736(20)32750-1.
- Muamala, T. (2020) 'Age categories according to WHO and Ministry of Health which are not widely known by the public'.
- Notoadmojo, S. (2012) *Promosi Kesehatan dan Perilaku Kesehatan*. Rineka Cipta.
- Putri, R. (2017). *The relationship between education and knowledge levels with healthy living behavior and the home environmental quality* (pp. 19–20). pp. 19–20. Retrieved from <http://digilib.unila.ac.id/26165/3/>.pdf
- Riyadi, R. and Larasaty, P. (2021) 'Faktor-Faktor yang Mempengaruhi Kepatuhan Masyarakat Terhadap Protokol Kesehatan dalam Mencegah Penyebaran Covid-19', *Seminar Nasional Official Statistics*, 2020(1), pp. 45–54. doi: 10.34123/semnasoffstat.v2020i1.431.
- Sardar, S. et al. (2022) 'Assessing the Effects of COVID-19 on Restaurant Business From Restaurant Owners ' Perspective', 13(April), pp. 1–10. doi: 10.3389/fpsyg.2022.849249.
- Semra Ozdemir , Sean Ng , Isha Chaudhry I, E. A. F. (2020). *Optimizing Health Protocol Enforcement during the Covid-19 Pandemic*. 16(2), 243–263. <https://doi.org/https://dx.doi.org/10.34172/ijhpm.2020.199>
- Sriarumtias, F. F. et al. (2020) 'Empowerment and Implementation of Health Protocols at Posyandu Leuwigoong Health Center, Garut Regency, West Java as an Effort Preventing the Transmission of COVID-19', *JURPIKAT (Jurnal Pengabdian Kepada Masyarakat)*, 1(1), pp. 1–12.
- Stewart, T., Day, S. W., Russell, J., Wilbanks, C., Likes, W., Webb, S., ... Cashion, A. K. (2020). Development of a COVID-19 alternate care site from ground zero: A nursing perspective. *Public Health Nursing*, 37(6), 889–894. <https://doi.org/10.1111/phn.12812>
- UNICEF (2020) *Important facts about hand washing with soap to protect you from coronavirus (COVID-19)*, Unicef Indonesia.
- Zandkarimi, M.;Ghobad;Behzad M. (2020) 'The Prognostic Factors Affecting the Survival of Kurdistan Province COVID-19 Patients: A Cross-sectional Study From February to May 2020', *International Journal of Health Policy and Management (IJHPM)*, x(x), pp. 1–6

How to cite this article: Erawati, A. D. (2022) 'Factors affecting compliance in implementing health protocols to prevent COVID-19 in Indonesia: A cross-sectional study', *Jurnal Ners*, 17(1), pp. 56-61. doi: <http://dx.doi.org/10.20473/jn.v17i1.25618>